


Order ORECTOLOBIFORMES

Nurse Sharks

Family GINGLYMOSOMATIDAE

Gill 1862



Nebrius concolor (= *ferrugineus*), type species of *Nebrius*. From: Rüppell, W. P. E. S. 1835–38. Neue Wirbelthiere zu der Fauna von Abyssinien gehörig. Fische des Rothen Meeres. Siegmund Schmerber, Frankfurt am Main. i–ii + 1–148, Pls. 1–33. [Pp. 53–80 and Pls. 15–21 published in 1837.]

Ginglymostoma

Müller & Henle 1837

ginglymos (Gr. γίγγλυμος), hinge; *stóma* (Gr. στόμα), mouth, presumably referring to how corner of mouth has a hinged appearance

Ginglymostoma cirratum (Bonnaterre 1788) Latin for having tendrils, referring to elongated nasal barbels

Ginglymostoma unami Del Moral-Flores, Ramírez-Antonio, Angulo & Pérez-Ponce de León 2015 in honor of UNAM (Universidad Nacional Autónoma de México), for the “vital role it plays in educating the people of Mexico” (translation)



Ginglymostoma unami, paratype, male, 207.5 cm TL. From: Del Moral-Flores, L. F., E. Ramírez-Antonio, A. Angula and G. Pérez-Ponce de León. 2015. *Ginglymostoma unami* sp. nov. (Chondrichthyes: Orectolobiformes: Ginglymostomidae): una especie nueva de tiburón gata del Pacífico oriental tropical. Revista Mexicana de Biodiversidad 86: 48–58.

Nebrius

Rüppell 1837

nebris (Gr. νεβρίς), fawn skin, referring to rusty-fawn coloration of upper body


Nebrius ferrugineus (Lesson 1831) Latin for rust-colored, referring to brown coloration

Pseudoginglymostoma

Dingerkus 1986

pseudo-, from *pseúdēs* (Gr. ψεύδης), false, i.e., not a true *Ginglymostoma*, in which *P. brevicaudatum* had originally been placed

Pseudoginglymostoma brevicaudatum (Günther 1867) *brevis* (L.), short; *caudatum* (L.), tailed, referring to caudal fin being less than $\frac{1}{4}$ of total length



Pseudoginglymostoma brevicaudatum. From: Playfair, R. L. and A. Günther. 1867. The fishes of Zanzibar, with a list of the fishes of the whole east coast of Africa. London. i–xiv + 1–153, Pls. 1–21.