

# Honey Composition

Stefan Bogdanov

## CHEMICAL COMPOSITION

The composition of honey depends mainly on the plant, from which the honey originates (see chapters Honey Elaboration and Honey Types) But it is influenced by different factors such as bee type, honey harvest, technology, harvest method, storage, soil type etc.


Honey is composed mainly from carbohydrates, lesser amounts of water and a great number of minor components.

**Honey composition** after <sup>76, 81</sup>, values in g/100 g


	Blossom honey		Honeydew honey	
	average	min-max	average	min-max
<b>Water content</b>	17.2	15-20	16.3	15-20
<b>Fructose</b>	38.2	30-45	31.8	28-40
<b>Glucose</b>	31.3	24-40	26.1	19-32
<b>Sucrose</b>	0.7	0.1-4.8	0.5	0.1-4.7
<b>Other disaccharides</b>	5.0	28	4.0	16
<b>Melezitose</b>	<0.1		4.0	0.3-22.0
<b>Erlose</b>	0.8	0.56	1.0	0.16
<b>Other oligosaccharides</b>	3.6	0.5-1	13.1	0.1-6
<b>Total sugars</b>	<b>79.7</b>		<b>80.5</b>	
<b>Minerals</b>	0.2	0.1-0.5	0.9	0.6-2
<b>Amino acids, proteins</b>	0.3	0.2-0.4	0.6	0.4-0.7
<b>Acids</b>	0.5	0.2-0.8	1.1	0.8-1.5
<b>pH</b>	3.9	3.5-4.5	5.2	4.5-6.5

**Further reading:** <sup>8, 44, 76</sup>


## Carbohydrates



**glucose**



**fructose**



**sucrose**

Sugars are the main constituents of honey, comprising about 95 % of honey dry weight. Main sugars are the monosaccharides hexoses fructose and glucose, which are products by the hydrolysis of the disaccharide sucrose. Besides, about 25 different sugars have been detected<sup>29, 62</sup>. The principal oligosaccharides in blossom honeys are disaccharides: sucrose, maltose, turanose, erlose. Honeydew honeys contain besides, also the trisaccharides melezitose and raffinose. Trace amounts of tetra and pentasaccharides have also been isolated.

The relative amount of the two monosaccharides fructose and glucose is useful for the classification of unifloral honeys<sup>12</sup>. On the other hand, the sugar spectra of minor sugars does not differ greatly in different blossom honeys<sup>12</sup>. This is due to the fact, that the oligosaccharides are mainly a product of honey invertase<sup>76</sup>. There are considerable differences between the sugar spectra of blossom and honeydew honeys, the latter containing a higher amount of oligosaccharides, mainly the trisaccharides melezitose and raffinose, both absent in blossom honeys (see table above) The differentiation between different types of honeydew honeys by sugar analysis is also difficult. An attempt to differentiate between honeydew honeys from various aphids was made by determination of specific oligosaccharides<sup>73</sup>. Metcalfa honey, a new honeydew honey type, produced mainly in Italy, can be distinguished from other honeydew honeys as it is rich in maltotriose and contains particularly high amounts of oligomers called dextrans<sup>36</sup>.

The sugar composition can be determined by different chromatographic methods<sup>11</sup>, HPLC being the most widely used one<sup>12</sup>.

**Further reading:** 19, 25, 47, 50

## Acidity and pH

The acid content of honey is relatively low but it is important for the honey taste. Most acids are added by the bees<sup>31</sup>. The main acid is gluconic acid, a product of glucose oxidation by glucose oxidase. However, it is present as its internal ester, a lactone, and does not contribute to honey's active acidity. Honey acidity is determined by titration<sup>11</sup> and is expressed in milli equivalents per kg. The following acid have been found in minor amounts: formic, acetic, citric, lactic, maleic, malic, oxalic, pyroglutamic and succinic<sup>48</sup>.

Most honeys are acidic, that means that the pH value is smaller than 7. The pH of blossom honeys varies between 3.3 to 4.6. An exception is the chestnut honey with a relatively high pH value of 5 to 6. Honeydew honeys, due to their higher mineral content, have a higher pH value, varying between 4.5 and 6.5. Honey is a buffer, that means that that its pH does not change by the addition of small quantities of acids and bases. The buffer capacity is due to the content of phosphates, carbonates and other mineral salts.

## Amino acids and proteins

The content of amino acids and proteins is relatively small, at the most 0.7 % (see table above).

Honey contains almost all physiologically important amino acids<sup>20, 53, 54</sup>. The main amino acid is proline is a measure of honey ripeness<sup>72</sup>. The proline content of normal honeys should be more than 200 mg/kg. Values below 180 mg/kg mean that the honey is probably adulterated by sugar addition.

The honey proteins are mainly enzymes, reviewed by White<sup>76</sup>. Bees add different enzymes during the process of honey ripening. Diastase (amylase) digests starch to maltose and is relatively stable to heat and storage. Invertase (saccharase,  $\alpha$ -glucosidase), catalyses mainly the conversion of sucrose to glucose and fructose, but also many other sugar conversions<sup>59</sup>. Two other main enzymes glucose oxidase and catalase regulate the production of  $H_2O_2$ , one of the honey antibacterial factors.


Diastase and invertase play an important role for judging of honey quality and are used as indicators of honey freshness. A minimum value of 8 diastase units is set in the Codex Alimentarius and the European honey directive. Their activity decay upon storage and heating of honey (see chapter 7). Invertase is more susceptible to damage by storage and heat and is used in some countries as an indicator for honey virginity and freshness. Fresh and virgin honeys are supposed to have at least 10 Hadorn invertase units<sup>30</sup>, or 64 International units, while honeys with low enzyme activity should have at least 4 units<sup>30, 74</sup>. The diastase and invertase activity vary in wide limits, depending on the botanical origin of honey<sup>55, 56</sup> and thus have a limited freshness indicating power. HMF is the better quality criterion in this respect.

**Further reading:** 2-5, 16, 38, 40, 42, 46, 51, 52, 54, 55, 57

## Hydroxymethylfurfuraldehyde (HMF)

Hydroxymethylfurfuraldehyde or HMF is a decomposition product of fructose. In fresh honey it is present only in trace amounts and its concentration increases with storage and prolonged heating of honey. The HMF building process depends on the pH, in the more acidic blossom honey it is built at a higher pace than in darker honeys with higher pH<sup>68</sup>. Short term heat treatment, even at higher temperature will increase the HMF content is only slightly<sup>65, 66</sup>, while upon storage of honey there is a steady temperature dependent increase of HMF. According to the Codex Alimentarius and EU standards the HMF maximum is 40 mg/kg, while honey from the tropics and blends with them should not have more than 80 mg/kg. Beekeeping organisations of some countries, e.g. Germany, Italy, Finland, Switzerland have set a maximum of 15 mg/kg for specially labelled “quality” or “virgin” honeys.

Building of HMF from a hexose sugar:



**Further reading:** 34, 39, 43, 58, 71, 75, 78

## Minerals und trace elements

Blossom honeys have a mineral content mostly between 0.1 and 0.3 %, while that of honeydew honeys can reach 1 % of the total. In early times the mineral content was determined as a quality criterion of honey. Today, this measurement is replaced by the determination of electrical conductivity.

Honey contains varying amounts of mineral substances ranging from 0.02 to 1.03 g/100 g<sup>76</sup>. The main element found in honey is potassium, besides many other elements (table...) Potassium, with an average of about one third of the total, is the main mineral element, but there is a wide variety of trace elements. Several investigations have shown that the trace element content of honey depends mainly on the botanical origin of honey, light blossom honeys having a lower content than dark honeys, e.g. honeydew, chestnut and heather<sup>35, 37, 61</sup>. It was possible to differentiate between different unifloral honeys by determination of different trace elements<sup>9, 49</sup>. The mineral and trace element content of honey was reviewed in 2016<sup>64</sup>

**Trace elements in honey**, after <sup>10, 64</sup>

Element	mg/100 g	Element	mg/100 g
Aluminium (Al)	0.01 - 2.4	Lead (Pb)*	0.001 - 0.03
Arsen (As)	0.014 - 0.026	Lithium (Li)	0.225 - 1.56
Barium (Ba)	0.01 - 0.08	Molybdenum (Mo)	0 - 0.004
Boron (B)	0.05 - 0.3	Nickel (Ni)	0 - 0.051
Bromine (Br)	0.4 - 1.3	Rubidium (Rb)	0.040 - 3.5
Cadmium (Cd)*	0 - 0.001	Silicium (Si)	0.05 - 24
Chlorine (Cl)	0.4 - 56	Strontium (Sr)	0.04 - 0.35
Cobalt (Co)	0.1 - 0.35	Sulfur (S)	0.7 - 26
Fluoride (F)	0.4 - 1.34	Vanadium (V)	0 - 0.013
Iodine (I)	10 - 100	Zirkonium (Zr)	0.05 - 0.08

\*- elements regarded as toxic, can be partially of anthropological origin

**Aroma compounds and phenolics**



Honey volatiles are the substances responsible for the honey aroma. Research on honey volatiles started in the early 1960s. Recently, by studying volatiles isolated from the blossom and from the respective unifloral honey, it was found that most volatile compounds originate probably from the plant, but some of them are added by bees <sup>1, 14</sup>. Until the present time about 600 compounds have been characterised in different honeys. As unifloral honeys differ in respect of their sensory properties, it is probable that analysis of volatile compounds will allow classification of unifloral honeys. Indeed, typical volatile substances have been found in many unifloral honey and analysis of volatile substances can be used for the authentication of the botanical origin of honey <sup>12, 21</sup>.

Phenolic acids and polyphenols are plant-derived secondary metabolites. These compounds have been used as chemotaxonomic markers in plant systematics. They have been suggested as possible markers for the determination of botanical origin of honey. Considerable differences in composition and content of phenolic compounds between different unifloral honeys were found. Dark coloured honeys are reported to contain more phenolic acid derivatives but less flavonoids than light coloured ones (Amiot et al., 1989). It was shown that most of the studied 9 European unifloral honeys can be distinguished by their typical flavonoid profile <sup>70</sup>. Honey samples contain also variable amounts of propolis-derived phenolic compounds that were not helpful for the determination of botanical origin. On the whole, the determination of the flavonoid patterns is useful for the classification of some but not all unifloral honeys. For a more in depth analysis of the flavonoid spectra of unifloral honeys see <sup>12, 28</sup>.

**Further reading:** <sup>12, 13, 21, 24, 26-28, 77</sup>

**Contaminants and toxic compounds**

Honey, as any other food can be contaminated from the environment, e.g. heavy metals, pesticides, antibiotics etc. <sup>7</sup>. Generally, the contamination levels found do not present a health hazard. The main problem today is contamination by antibiotics, used against the bee brood diseases. In the European Union antibiotics are not allowed to be used, and thus honey containing antibiotics is also not permitted on the market.

A few plants yield nectar containing toxic substances. There are two main toxin groups: diterpenoids and pyrazolidine alkaloids. Some plants of the *Ericaceae* family belonging to the sub-family *Rhododendron*, e.g. *Rhododendron ponticum* contain toxic polyhydroxylated cyclic hydrocarbons or diterpenoids <sup>22</sup>. Substance of the other toxin group, pyrazolidine alkaloids, are found in different honey types and the potential intoxication by these substances is reviewed <sup>32</sup>. Cases of honey poisoning have been reported very rarely in the literature and concern mostly individuals from following regions: Caucasus, Turkey, New Zealand, Australia, Japan, Nepal, South Africa and different countries in North and South America. The symptoms encountered after

honey poisoning are: vomiting, headache, stomach ache, unconsciousness, delirium, nausea, slight weakness. The poisonous plants are generally known to the beekeepers, thus honeys which can contain poisonous substances are not marketed. To minimise risks in countries where plants with poisonous nectar are growing, tourists are advised to buy honeys from the market and not from individual beekeepers.

The heavy metal content of honey was reviewed in 2016<sup>64</sup>

**Further reading:** <sup>7, 10</sup>

## MICROBIOLOGICAL COMPOSITION

### Bacteria

Honey, is a very concentrated sugar solution with a high osmotic pressure, making impossible the growth of any microorganisms. It contains less microorganisms than other natural food, especially there are no dangerous Bacillus species. Honey contains Bacillus bacteria, causing the dangerous bee pests, but these are not toxic for humans. That is why, in order to prevent bee pests, honey should not be disposed in open places, where it can easily be accessed by bees.

However, a number of bacteria are present in honey, most of them being harmless to man. Recent extensive reviews covered the main aspects of honey microbiology and the possible risks<sup>15, 63, 79</sup>.

The presence of *Clostridium botulinum* spores in honey was reported for the first time in 1976<sup>41</sup>. Since then there were many studies in honey all over the world. In some of them no *Botulinum* was found, in others, few honeys were found to contain the spores.<sup>15, 23, 63, 67</sup> Honey does not contain the *Botulinum* toxin, but the spores can theoretically build the toxin after digestion of infants until one year old. Very few cases of infant botulism after ingestion of honey have been reported lately and this has been attributed to *C. botulinum* spores present in honey. These findings have lead the health authorities of some countries (US, UK) to label honeys, that honey be not given to infants until one year of age. Most countries find that such notice is unnecessary. Indeed, honey is not the only source of *C. botulinum* spores as it can be found in any natural food.

In 2002 an expert study of the Health and Consumer Directorate of the European Commission carried out on "Honey and microbiological hazards"<sup>33</sup>. It was concluded that:

*“Although infant botulism is a serious illness, mortality is very low. In general, in Europe, the risk of infant botulism is extremely low. The majority of infants suffering from botulism have been given honey. The level and frequency of contamination of honey with spores of C. botulinum appear generally to be low, although limited microbiological testing of honey has been performed. The routes by which spores of C. botulinum contaminate honey have not been precisely identified.*

*Although some geographical regions of the world can be associated with a particular type of C. botulinum in the soil, it is not possible to identify countries as the origin of honey with a greater risk of containing C. botulinum.*

*C. botulinum can survive as spores in honey but cannot multiply or produce toxins due to the inhibitory properties of honey. At present there is no process that could be applied to remove or kill spores of C. botulinum in honey without impairing product quality.*

*Microbiological testing would not be an effective control option against infant botulism, due to the sporadic occurrence and low levels of C. botulinum in honey”.*

## Yeast

Honey contains naturally different osmotolerant yeast, which can cause undesirable fermentation. Osmotolerant yeasts can particularly develop in honeys with high moisture content.

In 1933 Lochhead<sup>45</sup> summarised investigations on the relationship of moisture content and fermentation on 319 honey samples as follows:



**Relationship of moisture content of honey and fermentation risk<sup>45</sup>**

Moisture content	Liability to fermentation
Less than 17.1	Safe regardless of yeast count
17.1-18 %	Safe if yeast count < 1000/g
18.1-19	Safe if yeast count < 10/g
19.1-20 %	Safe if yeast count < 1/g
Above 20 %	Always in danger

These conclusions, although based on old research, have been confirmed by practice. Some honey types, e.g. rape, sunflower and also honeys from tropical countries has a higher content of osmotolerant yeast and are less stable than other honeys with normal yeast counts<sup>69</sup>

Honey fermentation is undesirable. The easiest way to control is to harvest honey with low humidity. Also, it should be stored in air-tight vessels. Fermentation control is carried out by determination of yeast count, ethanol and glycerin content. Honey should comply to following quality criteria:

- Yeast count maximum 500000 per 10 g<sup>6, 60</sup>
- Glycerol, maximum content: 300 mg/kg<sup>6, 60</sup>
- Ethanol, maximum content 150 mg/kg<sup>80</sup>

**Further Reading :** 15, 17, 18, 63, 79

### References

1. ALISSANDRAKIS, E; DAFERERA, D; TARANTILIS, P A; POLISSIOU, M; HARIZANIS, P C (2003) Ultrasound-assisted extraction of volatile compounds from citrus flowers and citrus honey. *Food Chemistry* 82: 575-582.
2. ALONSO-TORRE, S R; CAVIA, M M; FERNANDEZ-MUINO, M A; MORENO, G; HUIDOBRO, J F; SANCHO, M T (2006) Evolution of acid phosphatase activity of honeys from different climates. *Food Chemistry* 97 (4): 750-755.
3. AZEREDO, L D; AZEREDO, M A A; DE SOUZA, S R; DUTRA, V M L (2003) Protein contents and physicochemical properties in honey samples of *Apis mellifera* of different floral origins. *Food Chemistry* 80 (2): 249-254.
4. BABACAN, S; PIVARNIK, L F; RAND, A G (2002) Honey amylase activity and food starch degradation. *Journal of Food Science* 67 (5): 1625-1630.
5. BABACAN, S; RAND, A G (2007) Characterization of honey amylase. *Journal of Food Science* 72 (1): C50-C55.
6. BECKH, G; LÜLLMANN, C (1999) Natürliche Bestandteile des Honigs - Hefen und deren Stoffwechselprodukte. Teil 1: Hefegehalt. *Deutsche Lebensmittel-Rundschau* 95 (11): 457-463.
7. BOGDANOV, S (2006) Contaminants of bee products. *Apidologie* 38 (1): 1-18.



## The Honey Book, Chapter 5

8. BOGDANOV, S; BIERI, K; FIGAR, M; FIGUEIREDO, V; IFF, D; KÄNZIG, A; STÖCKLI, H; ZÜRCHER, K (1995) Kapitel 23 Bienenprodukte: 23A Honig. *Schweiz.Lebensmittelbuch* (11)
9. BOGDANOV, S; HALDIMANN, M; LUGINBÜHL, W; GALLMANN, P (2007) Minerals in honey: environmental, geographical and botanical aspects. *Journal of Apicultural Research and Bee World* 46 (4): 269-275.
10. BOGDANOV, S; JURENDIC, T; SIEBER, R; GALLMANN, P (2008) Honey for Nutrition and Health: A Review. *J.Am..Coll.Nutr.* 27: 677-689.
11. BOGDANOV, S; MARTIN, P; LÜLLMANN, C (1997) Harmonised methods of the European honey commission. *Apidologie* (extra issue): 1-59.
12. BOGDANOV, S; RUOFF, K; PERSANO ODDO, L (2004) Physico-chemical methods for the characterisation of unifloral honeys: a review. *Apidologie* 35 (Special issue): 4-17.
13. BONVEHI, J S; COLL, F V (2003) Flavour index and aroma profiles of fresh and processed honeys. *Journal of the Science of Food and Agriculture* 83 (4): 275-282.
14. CEPURNOI, I (2002) *Expertise in honey quality*. Editing House Dashkov and company, Moscow Russia
15. CLIVER, D O (2000) Honey, human pathogens, and HACCP. *Dairy, Food and Environmental Sanitation* 20 (4): 261-263.
16. COMETTO, P M; FAYE, P F; CACCAVARI, M; BARONI, M V; ALDAO, M A J (2006) Relationship between interannual variation of amino acid profile and pollen content in honey from a small Argentinian region. *Journal of agricultural and food chemistry* 54 (25): 9458-9464.
17. COMI, G; MANZANO, M; LENARDON, M; COCOLIN, L; CANTONI, C (2000) Evaluation of the parameters (a(w), humidity, storage temperature and osmophilic yeasts concentration) influencing yeast fermentation in honey. *Industria Alimentari* 39 (396): 1127-1133.
18. COMI, G; MANZANO, M; LENARDON, M; COCOLIN, L; GAIDELLA, L (2000) Microbiological and chemical-physical aspects of various honeys. *Industria Alimentari* 39 (395): 966-975.
19. COTTE, J F; CASABIANCA, H; CHARDON, S; LHERITIER, J; GRENIER-LOUSTALOT, M F (2003) Application of carbohydrate analysis to verify honey authenticity. *Journal of Chromatography.A* 1021 (1-2): 145-155.
20. COTTE, J F; CASABIANCA, H; GIROUD, B; ALBERT, M; LHERITIER, J; GRENIER-LOUSTALOT, M F (2004) Characterization of honey amino acid profiles using high- pressure liquid chromatography to control authenticity. *Analytical and Bioanalytical Chemistry* 378 (5): 1342-1350.
21. CUEVAS-GLORY, L F; PINO, J A; SANTIAGO, L S; SAURI-DUCH, E (2007) A review of volatile analytical methods for determining the botanical origin of honey. *Food Chemistry* 103 (3): 1032-1043.
22. DE BODT, G (1996) Les miels de rhododendrons. *Les Carnets du CARI Abeilles et Cie* (50): 10-12.
23. DE BODT, G; VLAYEN, P (1994) Miel et botulisme. *Les Carnets du CARI Abeilles et Cie* (46): 14-16.
24. DE LA FUENTE, E; MARTINEZ-CASTRO, I; SANZ, J (2005) Characterization of Spanish unifloral honeys by solid phase microextraction and gas chromatography-mass spectrometry. *Journal of Separation Science* 28 (9-10): 1093-1100.
25. DE LA FUENTE, E; SANZ, M L; MARTINEZ-CASTRO, I; SANZ, J (2006) Development of a robust method for the quantitative determination of disaccharides in honey by gas chromatography. *Journal of Chromatography.A* 1135 (2): 212-218.
26. DE LA FUENTE, E; SANZ, M L; MARTINEZ-CASTRO, I; SANZ, J; RUIZ-MATUTE, A I (2007) Volatile and carbohydrate composition of rare unifloral honeys from Spain. *Food Chemistry* 105 (1): 84-93.

## The Honey Book, Chapter 5

27. DE MARIS, C A B; MOREIRA, R F A (2003) Volatile compounds in floral honeys. *Quimica Nova* 26 (1): 90-96.
28. DIMITROVA, B; GEVRENOVA, R; ANKLAM, E (2007) Analysis of phenolic acids in honeys of different floral origin by solid-phase extraction and high-performance liquid chromatography. *Phytochemical Analysis* 18 (1): 24-32.
29. DONER, L W (1977) The sugars of honey - a review. *Journal of the Science of Food and Agriculture* 28: 443-456.
30. DUISBERG, H; HADORN, H (1966) Welche Anforderungen sind an Handelshonige zu stellen? *Mitteilungen aus dem Gebiete der Lebensmitteluntersuchung und Hygiene* 57: 386-407.
31. ECHIGO, T; TAKENAKA, T (1974) Production of organic acids in honey by honeybees. *J.Agric.Chem.Soc.Japan* 48 (4): 225-230.
32. EDGAR, J A; ROEDER, E L; MOLYNEUX, R J (2002) Honey from plants containing pyrrolizidine alkaloids: A potential threat to health. *Journal of agricultural and food chemistry* 50 (10): 2719-2730.
33. EUROPEAN COMMISSION (2002) Honey and microbiological hazards. *Report European Commission of Health & Consumer Protection Directorate-General*: 1-40.
34. FALLICO, B; ZAPPALA, M; ARENA, E; VERZERA, A (2004) Effects of conditioning on HMF content in unifloral honeys. *Food Chemistry* 85 (2): 305-313.
35. FELLER-DEMALSY, M J; VINCENT, B; BEAULIEU, F (1989) Mineral content and geographical origin of Canadian honeys. *Apidologie* 20 (1): 77-91.
36. FIORI, J; SERRA, G; SABATINI, A G; ZUCCHI, P; BARBATTINI, R; GAZZIOLA, F (2000) Dextrins HPLC analysis in *Metcalfa pruinosa* (Say) honeydew. *Industria Alimentari* 39 (391): 463-466.
37. GONZALEZ-MIRET, M L; TERRAB, A; HERNANZ, D; FERNANDEZ-RECAMALES, M A; HEREDIA, F J (2005) Multivariate correlation between color and mineral composition of honeys and by their botanical origin. *Journal of agricultural and food chemistry* 53 (7): 2574-2580.
38. GONZALEZ-PARAMAS, A M; GARCIA-VILLANOVA, R J; BAREZ, J A G; SANCHEZ, J S; ALBAJAR, R A (2007) Botanical origin of monovarietal dark honeys (from heather, holm oak, pyrenean oak and sweet chestnut) based on their chromatic characters and amino acid profiles. *European food research and technology* 226 (1-2): 87-92.
39. HEBBAR, H U; NANDINI, K E; LAKSHMI, M C; SUBRAMANIAN, R (2003) Microwave and infrared heat processing of honey and its quality. *Food Science and Technology Research* 9 (1): 49-53.
40. HERMOSIN, I; CHICON, R M; CABEZUDO, M D (2003) Free amino acid composition and botanical origin of honey. *Food Chemistry* 83 (2): 263-268.
41. HUHTANEN, C N; KNOX, D; SHIMANUKI, H (1981) Incidence and origin of *Clostridium botulinum* spores in honey. *Journal of Food Protection* 44 (11): 812-814.
42. IGLESIAS, M T; MARTIN-ALVAREZ, P J; POLO, M C; DE LORENZO, C; GONZALEZ, M; PUEYO, E (2006) Changes in the free amino acid contents of honeys during storage at ambient temperature. *Journal of agricultural and food chemistry* 54 (24): 9099-9104.
43. KUBIS, I; INGR, I (1998) Effects inducing changes in hydroxymethylfurfural content in honey. *Czech Journal of Animal Science* 43 (8): 379-383.
44. LIPP, J; ZANDER, E; KOCH, A (1994) *Der Honig*. Eugen Ulmer Stuttgart
45. LOCHHEAD, A G (1933) Factors concerned with the fermentation of honey. *Zent.Bakt.Paras.u.Infect.II* 88: 296-302.



## The Honey Book, Chapter 5

46. LOW, N H; VONG, K V; SPORNS, P (1986) A new enzyme,  $\alpha$ -glucosidase, in honey. *Journal of Apicultural Research* 25 (3): 178-181.
47. MATEO, R; BOSCH-REIG, F (1997) Sugar profiles of Spanish unifloral honeys. *Food Chemistry* 60 (1): 33-41.
48. MATO, I; HUIDOBRO, J F; SIMAL-LOZANO, J; SANCHO, M T (2003) Significance of nonaromatic organic acids in honey. *Journal of Food Protection* 66 (12): 2371-2376.
49. NOZAL NALDA, M J; YAGUE, J L B; CALVA, J C D; GOMEZ, M T M (2005) Classifying honeys from the Soria Province of Spain via multivariate analysis. *Analytical and Bioanalytical Chemistry* 382 (2): 311-319.
50. NOZAL, M J; BERNAL, J L; TORIBIO, L; ALAMO, M; DIEGO, J C; TAPIA, J (2005) The use of carbohydrate profiles and chemometrics in the characterization of natural honeys of identical geographical origin. *Journal of agricultural and food chemistry* 53 (8): 3095-3100.
51. ODDO, L P; PIAZZA, M G; PULCINI, P (1999) Invertase activity in honey. *Apidologie* 30 (1): 57-65.
52. PAWLOWSKA, M; ARMSTRONG, D W (1994) Evaluation of enantiomeric purity of selected amino acids in honey. *Chirality* 6: 270-276.
53. PEREZ, C; CONCHELLO, P; ARINO, A; YANGUELA, J; HERRERA, A (1989) Dosage des acides aminés des protéines de différents miels espagnols. *Sciences des Aliments* 9: 203-207.
54. PEREZ, R A; IGLESIAS, M T; PUEYO, E; GONZALEZ, M; DE LORENZO, C (2007) Amino acid composition and antioxidant capacity of Spanish honeys. *Journal of agricultural and food chemistry* 55 (2): 360-365.
55. PERSANO ODDO, L; BALDI, E; ACCORTI, M (1990) Diastatic activity in some unifloral honeys. *Apidologie* 21 (1): 17-24.
56. PERSANO ODDO, L; PIAZZA, M G; PULCINI, P (1999) Invertase activity in honey. *Apidologie* 30 (1): 57-65.
57. QAMER, S; EHSAN, M; NADEEM, S; SHAKOORI, A R (2007) Free amino acids content of Pakistani unifloral honey produced by *Apis mellifera*. *Pakistan Journal of Zoology* 39 (2): 99-102.
58. RAMIREZ CERVANTES, M A; GONZALES NOVELO, S A; SAURI DUCH, E (2000) Die Wirkung der zeitweiligen Wärmebehandlung des Honigs auf seine qualitativen Variationen während der Lagerung. *Apiacta* 35 (4): 162-170.
59. RAUDE-ROBERG, L (1994) Analyse der Saccharide Spaltenden Enzyme des Bienenhonigs. Dr. Dissertation; Celle, Deutschland Niedersächsisches Landesinstitut für Bienenkunde
60. RUSSMANN, H (1998) Hefen und Glycerin in Blütenhonigen - Nachweis einer Gärung oder einer abgestoppten Gärung. *Lebensmittelchemie* 52: 116-117.
61. SEVLIMLI, H; BAYULGEN, N; VARINIOGLU (1992) Determination of trace elements in honey by INAA in Turkey. *J.Radioanal.Nucl.Chem., Letters* 165 (5): 319-325.
62. SIDDIQUI, I R (1970) The sugars of honey. *Advances in Carbohydrate Chemistry and Biochemistry* 25: 285-309.
63. SNOWDON, J A; CLIVER, D O (1996) Microorganisms in honey. *International Journal of Food Microbiology* 31 (1/3): 1-26.
64. SOLAYMAN, M I M P S A Y K M A N & G S H (2016) Physicochemical Properties, Minerals, Trace Elements, and Heavy Metals in Honey of Different Origins: A Comprehensive Review. *Comprehensive Reviews in Food Science and Food Safety*, 15: 219-233.

## The Honey Book, Chapter 5

65. SUBRAMANIAN, R; HEBBAR, H U; RASTOGI, N K (2007) Processing of honey: A review. *INTERNATIONAL JOURNAL OF FOOD PROPERTIES* 10 (1): 127-143.
66. TABOURET, T; MATHLOUTHI, M (1972) Essai de pasteurisation de miel. *Rev.franc.Apic.* 299: 258-261.
67. TANZI, M G; GABAY, M P (2002) Association between honey consumption and infant botulism. *Pharmacotherapy* 22 (11): 1479-1483.
68. THRASYVOULOU, A (1997) Heating times for Greek honeys. *Melissokomiki Epitheorisi* 11 (2): 79-80.
69. TIMMROTH, R; SPEER, K; BECKH, G; LÜLLMANN, C (2005) Comparison of European honeys to tropical honeys - effects of yeast cell numbers on the concentration of especially selected components *Apimondia abstracts Ireland 2005*, Apimondia International Apicultural Congress Dublin, Ireland; Dublin, Ireland; pp 110.
70. TOMÁS-BARBERÁN, F A; MARTOS, I; FERRERES, F; RADOVIC, B S; ANKLAM, E (2001) HPLC flavonoid profiles as markers for the botanical origin of European unifloral honeys. *Journal of the Science of Food and Agriculture* 81 (5): 485-496.
71. TOSI, E; CIAPPINI, M; RE, E; LUCERO, H (2002) Honey thermal treatment effects on hydroxymethylfurfural content. *Food Chemistry* 77 (1): 71-74.
72. VON DER OHE, W; DUSTMANN, J H; VON DER OHE, K (1991) Prolin als Kriterium der Reife des Honigs. *Deutsche Lebensmittel-Rundschau* 87 (12): 383-386.
73. VON DER OHE, W; VON DER OHE, K (1996) Characterisation of honeydew honey guided with specific saccharides. Charakterisierung von Honigtauhonig anhand spezifischer Saccharide. *Apidologie* 27 (4): 270-272.
74. VON DER OHE, W; VON DER OHE, K; RAUDE-ROBERG, L; DUSTMANN, J H (1999) Vergleich der Methoden zur Bestimmung der Saccharase-Aktivität im Honig. *Apidologie* 30: 412-413.
75. WEIGEL, K U; OPITZ, T; HENLE, T (2004) Studies on the occurrence and formation of 1,2-dicarbonyls in honey. *European food research and technology* 218 (2): 147-151.
76. WHITE, J W (1975) Composition of honey., In Crane, E (ed.) *Honey, a comprehensive survey*, Heinemann Edition; London; pp 157-206.
77. WOLSKI, T; TAMBOR, K; RYBAK-CHMIELEWSKA, H; KEDZIA, B (2006) Identification of honey volatile components by solid phase microextraction (SPME) and gas chromatography/mass spectrometry (GC/MS). *Journal of Apicultural Science* 50 (2): 115-126.
78. WUNDERLIN, D A; PESCE, S F; AME, M V; FAYE, P F (1998) Decomposition of hydroxymethylfurfural in solution and protective effect of fructose. *Journal of agricultural and food chemistry* 46 (5): 1855-1863.
79. ZUCCHI, P; BASSIGNANI, V; CARPANA, E (2001) Honey microbiology. *Industria Alimentari* 40 (409): 1346-1350.
80. ZUCCHI, P; MARCAZZAN, G L; DAL POZZO, M; SABATINI, A G; DESALVO, F; FLORIS, I (2006) Il contenuto di etanolo nel miele per la valutazione di processi fermentativi. *APOidea* 3 (1): 18-26.
81. [ANON] (1995) *Swiss food manual, Chapter 23 A, Honey*. Eidgenössische Druck und Materialzentrale Bern