



PRODUCT DESCRIPTION

RofenAid[®] 40 is a Type A Medicated Article for the prevention of coccidiosis and bacterial infections in poultry. RofenAid is an antibacterial anticoccidial Type A Medicated Article containing 113.4 grams sulfadimethoxine (25%) and 68.1 grams ormetoprim (15%) per pound of premix.

INDICATIONS FOR USE

- Aids in the prevention of coccidiosis in chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*; as an aid in the prevention of bacterial infections caused by *Haemophilus gallinarum* (infectious Coryza), *Escherichia coli* (colibacillosis) and *Pasteurella multocida* (fowl cholera)
- Aids in the prevention of coccidiosis in turkeys caused by *Eimeria adenoides*, *E. gallopavonis* and *E. meleagrimitis*; for the prevention of bacterial infections caused by *Pasteurella multocida* (fowl cholera)
- Aids in the control of *Pasteurella multocida* (fowl cholera) in ducks and also as an aid in the control of *E. coli* and *Riemerella anatipestifer*
- For the prevention of coccidiosis in chukar partridges caused by *Eimeria kofoidi* and *E. legionensis*

KEY POINTS

- Antibacterial and anticoccidial combination
- The only sulfonamide potentiated with ormetoprim for enhanced broad-spectrum antibacterial activity, compared with sulfonamides alone¹

FDA STATUS

- Feeds containing RofenAid 40 should be fed as the sole ration
- Category II drug and a feed mill license is required

STORAGE AND PACKAGING

- Store below 25° C (77° F), excursions permitted to 37° C (99° F)
- RofenAid 40 is packaged in a 50-pound multiwall paper bag with protective barrier ply
- Expiration period is 24 months from the date of manufacture (package indicates expiration date)

WITHDRAWAL INFORMATION

- Do not feed to chickens over 16 weeks of age
- Do not feed to turkeys and ducks producing eggs for human consumption
- A 5-day withdrawal is required prior to slaughter of broiler and replacement chickens, turkeys and ducks

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Sulfadimethoxine and Ormetoprim — Type A Medicated Article

Active Drug Ingredients	Sulfadimethoxine.....113.5 grams per pound (25%) Ormetoprim..... 68.1 grams per pound (15%) in a carrier suitable for incorporation in feed.
For use in manufacturing medicated feeds only	
Broiler and replacement (breeder and layer) chickens	INDICATIONS: As an aid in the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> and <i>E. mivati</i> ; as an aid in the prevention of bacterial infections caused by <i>Haemophilus gallinarum</i> (infectious Coryza), <i>Escherichia coli</i> (colibacillosis) and <i>Pasteurella multocida</i> (fowl cholera). USE DIRECTIONS: Add 1 lb. of RofenAid 40 to 1 ton of feed to provide 0.02% (113.5 g/ton [0.0125%] sulfadimethoxine and 68.1 g/ton [0.0075%] ormetoprim) concentration in feed. Feed continuously as the sole ration. WARNING: Do not feed to chickens over 16 weeks of age (112 days).
Turkeys	INDICATIONS: As an aid in the prevention of coccidiosis caused by <i>Eimeria adenoeides</i> , <i>E. gallopavonis</i> and <i>E. meleagriditis</i> ; for the prevention of bacterial infections caused by <i>Pasteurella multocida</i> (fowl cholera). USE DIRECTIONS: Add 0.5 lb. of RofenAid 40 to 1 ton of feed to provide 0.01% (56.75 g/ton [0.00625%] sulfadimethoxine and 34.05 g/ton [0.00375%] ormetoprim) concentration in feed. Feed continuously as the sole ration.
Chukar partridges	INDICATIONS: For the prevention of coccidiosis caused by <i>Eimeria kofoidi</i> and <i>E. legionensis</i> . USE DIRECTIONS: Add 1 lb. of RofenAid 40 to 1 ton of feed to provide 0.02% (113.5 g/ton [0.0125%] sulfadimethoxine and 68.1 g/ton [0.0075%] ormetoprim) concentration in feed. Feed continuously as the sole ration to young birds up to 8 weeks of age. CAUTION: The safety of feeding RofenAid 40 to breeding stock has not been established.
Breeding ducks	INDICATIONS: As an aid in the control of bacterial infections due to <i>Pasteurella multocida</i> (fowl cholera) in ducks, including breeding ducks. USE DIRECTIONS: Add 2 lbs. of RofenAid 40 to 1 ton of feed to provide 0.04% (227 g/ton [0.025%] sulfadimethoxine and 136.2 g/ton [0.015%] ormetoprim) concentration in feed. Medication should be started at the first signs of infection. Feed for 7 days as the sole ration.
Ducks	INDICATIONS: As an aid in the control of bacterial infections due to <i>Escherichia coli</i> , <i>Riemerella anatipestifer</i> (strains 2, 3 & 4) and severe challenge of <i>Pasteurella multocida</i> (fowl cholera). USE DIRECTIONS: Add 4 lbs of RofenAid 40 to 1 ton of feed to provide 0.08% (454 g/ton [0.050%] sulfadimethoxine and 272.4 g/ton [0.030%] ormetoprim) concentration in feed. Medication should be started at the first signs of infection. Feed for 7 days as the sole ration. WARNING: Not for breeding ducks. The safety of feeding sulfadimethoxine and ormetoprim at levels higher than 0.025% and 0.015%, respectively, in breeding ducks has not been established.
Warning	Do not feed to turkeys and ducks producing eggs for food. Withdraw 5 days prior to slaughter of broiler and replacement chickens, turkeys and ducks.

NOTE: Manufacture of Type B or C feeds from this product requires a Medicated Feed Mill License

Labels for feeds containing RofenAid 40 must contain appropriate indications, limitations and warnings as well as required feed ingredient information.

Restricted Drug (California) - Use only as directed. Not for human use.

Store below 25° C (77° F), excursions permitted to 37° C (99° F)

Approved by FDA under NADA #040-209

¹McDougald LR, Roberson EL. Antiprotozoan drugs. In: Booth NH, McDonald LE, eds. *Veterinary Pharmacology and Therapeutics*. Ames, Iowa: Iowa State Press, 1988;950-968.

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted.
© 2020 Zoetis Services LLC. All rights reserved. PHA-00059R1