

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Albumin Octapharma 200 g/l
Human albumin 200 g/l
Solution for infusion

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Human albumin 200 g/l*

*solution containing 200g/l of proteins of which at least 96% is human albumin;

One bottle of 50 ml contains 10 g of human albumin.

One bottle of 100 ml contains 20 g of human albumin.

Albumin 200 g/l is a hyperoncotic solution.

Excipients:

Sodium 142.5-157.5 mmol/l

Potassium max. 2.0 mmol/l

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Solution for infusion.

Clear or slightly opalescent.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Restoration and maintenance of circulating blood volume where volume deficiency has been demonstrated, and use of a colloid is appropriate.

The choice of albumin rather than artificial colloid will depend on the clinical situation of the individual patient, based on official recommendations.

4.2 Posology and method of administration

The concentration of the albumin preparation, dosage and the infusion-rate should be adjusted to the patient's individual requirements.

Posology

The dose required depends on the size of the patient, the severity of trauma or illness and on continuing fluid and protein losses. Measures of adequacy of circulating volume and not plasma albumin levels should be used to determine the dose required.

If human albumin is to be administered, haemodynamic performance should be monitored regularly; this may include:

- arterial blood pressure and pulse rate
- central venous pressure
- pulmonary artery wedge pressure
- urine output
- electrolyte
- haematocrit/haemoglobin

This product is suitable for premature infants and dialysis patients.

Method of administration

Human albumin can be directly administered by the intravenous route.

The infusion rate should be adjusted according to the individual circumstances and the indication.

In plasma exchange the infusion rate may be higher and should be adjusted to the rate of removal.

4.3 Contraindications

Hypersensitivity to albumin preparations or to any of the excipients.

4.4 Special warnings and precautions for use

If allergic or anaphylactic-type reactions occur, the infusion should be stopped immediately and appropriate treatment instituted. In case of shock, the current medical standards for shock-treatment should be observed.

Albumin should be used with caution in conditions where hypervolaemia and its consequences or haemodilution could represent a special risk for the patient. Examples of such conditions are:

- Decompensated cardiac insufficiency
- Hypertension
- Oesophageal varices
- Pulmonary oedema
- Haemorrhagic diathesis
- Severe anaemia
- Renal and post-renal anuria

The colloid-osmotic effect of human albumin 200 g/l is approximately four times that of blood plasma. Therefore, when concentrated albumin is administered, care must be taken to assure adequate hydration of the patient. Patients should be monitored carefully to guard against circulatory overload and hyperhydration.

20-25% human albumin solutions are relatively low in electrolytes compared to the 4-5% human albumin solutions. When albumin is given the electrolyte status of the patient should be monitored (see 4.2) and appropriate steps taken to restore or maintain the electrolyte balance.

Albumin solutions must not be diluted with water for injections as this may cause haemolysis in recipients.

If comparatively large volumes are to be replaced, controls of coagulation and haematocrit are necessary. Care must be taken to ensure adequate substitution of other blood constituents (coagulation factors, electrolytes, platelets and erythrocytes).

Hypervolaemia may occur if the dosage and rate of infusion are not adjusted to the patients circulatory situation. At the first clinical signs of cardiovascular overload (headache, dyspnoea, jugular vein congestion), or increased blood pressure, raised venous pressure and pulmonary oedema, the infusion is to be stopped immediately.

This medicinal product contains 14.25 –15.75 mmol sodium per 100 ml solution. This has to be taken into consideration by patients on a controlled sodium diet.

This medicine contains less than 1 mmol potassium per 100 ml solution, i.e. essentially “potassium-free”.

Standard measure to prevent infections resulting from the use of medicinal products prepared from human blood or plasma include selection of donors, screening of individual donations and plasma pools for specific markers of infection and the inclusion of effective manufacturing steps for the inactivation/removal of viruses. Despite this, when medicinal products prepared from human blood or plasma are administered, the possibility of transmitting infective agents cannot be totally excluded. This also applies to unknown or emerging viruses and other pathogens.

There are no reports of virus transmissions with albumin manufactured to European Pharmacopeia specifications by established processes.

It is strongly recommended that every time that Albumin Octapharma is administered to a patient, the name and batch number of the product are recorded in order to maintain a link between the patient and the batch of the product.

4.5 Interactions with other medicinal products and other forms of interactions

No specific interactions of human albumin with other medicinal products are known.

4.6 Pregnancy and lactation

The safety of Albumin Octapharma for use in human pregnancy has not been established in controlled clinical trials. However, clinical experience with albumin suggests that no harmful effects on the course of pregnancy, or on the foetus and the neonate are to be expected.

No animal reproduction studies have been conducted with Albumin Octapharma. However, human albumin is a normal constituent of human blood.

4.7 Effects on ability to drive and use machines

No effects on ability to drive and use machines have been observed.

4.8 Undesirable effects

Adverse reactions for Albumin Octapharma are rare. These reactions normally disappear rapidly when the infusion rate is slowed down or the infusion is stopped. In case of severe reactions, the infusion should be stopped and an appropriate treatment should be initiated.

The following adverse reactions have been observed for Albumin Octapharma during the postmarketing phase.

Very common (>1/10); common (>1/100, <1/10); uncommon (>1/1,000, <1/100); rare (>1/10,000, <1/1,000); very rare (<1/10,000), including isolated reports.

System Organ Class	Rare	Very rare
<i>Immune system disorders</i>	anaphylactic reaction hypersensitivity	anaphylactic shock
<i>Psychiatric disorders</i>		confusional state
<i>Nervous system disorders</i>		headache
<i>Cardiac disorders</i>		tachycardia bradycardia
<i>Vascular disorders</i>	hypotension	hypertension flushing
<i>Respiratory, thoracic and mediastinal disorders</i>		dyspnoea
<i>Gastrointestinal disorders</i>		nausea
<i>Skin and subcutaneous tissue disorders</i>		urticaria angioneurotic oedema rash erythematous hyperhidrosis
<i>General disorders and</i>		pyrexia

System Organ Class	Rare	Very rare
<i>administration site conditions</i>		chills

For information on viral safety see 4.4.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V*.

4.9 Overdose

Hypervolaemia may occur if the dosage and rate of infusion are too high. At the first clinical signs of cardiovascular overload (headache, dyspnoea, jugular vein congestion), or increased blood pressure, raised central venous pressure and pulmonary oedema, the infusion should be stopped immediately and the patient's haemodynamic parameters carefully monitored.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: plasma substitutes and plasma protein fractions, ATC code: B05AA01

Human albumin accounts quantitatively for more than half of the total protein in the plasma and represents about 10% of the protein synthesis activity of the liver.

Physiochemical data:

Human albumin 200 g/l has a corresponding hyperoncotic effect.

The most important physiological function of albumin results from its contribution to oncotic pressure of the blood and transport function. Albumin stabilises circulating blood volume and is a carrier of hormones, enzymes, medicinal products and toxins.

5.2 Pharmacokinetic properties

Under normal conditions the total exchangeable albumin pool is 4-5 g/kg body weight of which 40-45% is present intravascularly and 55-60% in the extravascular space. Increased capillary permeability will alter albumin kinetics and abnormal distribution may occur in conditions such as severe burns or septic shock.

Under normal conditions, the average half-life of albumin is about 19 days. The balance between synthesis and breakdown is normally achieved by feed-back regulation. Elimination is predominantly intracellular and due to lysosome proteases.

In healthy subjects, less than 10% of infused albumin leaves the intravascular compartment during the first 2 hours following infusion. There is considerable individual variation in the effect on plasma volume. In some patients the plasma volume can remain increased for some hours. However, in critically ill patients, albumin can leak out of the vascular space in substantial amounts at an unpredictable rate.

5.3 Preclinical safety data

Human albumin is a normal constituent of human plasma and acts like physiological albumin.

In animals, single dose toxicity testing is of little relevance and does not permit the evaluation of toxic or lethal doses or of a dose-effect-relationship. Repeated dose toxicity testing is impracticable due to the development of antibodies to heterologous protein in animal models.

To date, human albumin has not been reported to be associated with embryo-foetal toxicity, oncogenic or mutagenic potential.

No signs of acute toxicity have been described in animal models.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

N-acetyl-DL-tryptophan	max. 16.8 mmol/l
Caprylic acid	max. 16.8 mmol/l
Water for injections	ad 1000 ml

Electrolytes

Sodium*	142.5-157.5	mmol/l
Potassium	max. 2.0	mmol/l

*(mainly from sodium chloride and sodium hydroxide)

6.2 Incompatibilities

Human albumin solution must not be mixed with other medicinal products, whole blood and packed red cells.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store and transport below 25 °C.

Keep the container in the outer carton in order to protect from light.

Do not freeze.

6.5 Nature and contents of container

- 50 ml of solution in infusion bottle (type II glass) with stopper (bromobutyl rubber). Pack of 1.
- 100 ml of solution in infusion bottle (type II glass) with stopper (bromobutyl rubber). Pack of 1.

6.6 Special precautions for disposal and other handling

The solution can be directly administered by the intravenous route.

Albumin solutions must not be diluted with water for injections as this may cause haemolysis in recipients.

If large volumes are administered, the product should be warmed to room temperature before use.

The solution should be clear or slightly opalescent. Do not use solutions which are cloudy or have deposits. This may indicate that the protein is unstable or that the solution has become contaminated.

Once the infusion container has been opened the content should be used immediately.

Any unused product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

[to be completed nationally]

8 MARKETING AUTHORISATION NUMBER(S)

[to be completed nationally]

9 DATE OF FIRST AUTHORISATION/ RENEWAL OF THE AUTHORISATION

1999-11-04/2009-11-04

10 DATE OF REVISION OF THE TEXT

27 March, 2015