

Aerosolized Prostacyclins

Mark Siobal RRT

Introduction

Prostacyclins Available for Aerosolization

Prostaglandin I₂

Iloprost

Prostaglandin E₁

Indications and Evidence for Aerosolized Prostacyclin

Pulmonary Hypertension

Right Ventricular Failure

Acute Respiratory Distress Syndrome

Other Potential Benefits

Adverse Effects and Toxicity

Aerosol Delivery Methods

Efficacy of Aerosolized Prostacyclin

Summary

Two prostacyclins (prostaglandin E₁ and prostaglandin I₂) are potent vasodilators. Aerosolized prostacyclins reduce pulmonary artery pressure, improve right heart function, and increase arterial oxygenation by improving ventilation/perfusion matching. This report describes aerosolized prostacyclins and compares them to inhaled nitric oxide. I review the types of inhalable prostacyclins and their indications, evidence of efficacy, delivery, and adverse effects. Key words: aerosol, prostacyclin, prostaglandin, pulmonary hypertension, ventricular function, acute respiratory distress syndrome, ARDS, inhaled nitric oxide, nebulizer. [Respir Care 2004;49(6):640–652. © 2004 Daedalus Enterprises]

Introduction

Inhaled vasodilators can reduce pulmonary artery pressure and redistribute pulmonary blood flow to ventilated lung regions, with little or no systemic hemodynamic effect.¹ The potential benefits of such targeted pulmonary

vasodilation include reduced pulmonary vascular resistance (PVR), reduced right ventricular afterload, improved right-heart function, better ventilation/perfusion (\dot{V}/\dot{Q}) matching, and improved arterial oxygenation (Fig. 1).²

The utility of inhaled prostacyclins was explored several years before the identification of nitric oxide^{3–5} as endothelium-derived relaxing factor.⁶ During the 1990s there was extensive research on inhaled nitric oxide (INO) as a vasodilator, in both animals and humans. INO significantly reduces the need for extracorporeal membrane oxygenation among near-term neonates who require mechanical ventilation, and so INO was approved by the Food and Drug Administration (FDA) for treatment of persistent pulmonary hypertension and hypoxemia.

In December 2000 San Francisco General Hospital's pharmacy and therapeutics committee added INO to the

Mark Siobal RRT is affiliated with Respiratory Care Services, Department of Anesthesia, University of California, San Francisco, at San Francisco General Hospital, San Francisco, California.

Mark Siobal RRT presented a version of this report at the 49th International Respiratory Congress, held December 8–11, 2003, in Las Vegas, Nevada.

Correspondence: Mark Siobal RRT, Respiratory Care Services, San Francisco General Hospital, NH GA-2, 1001 Potrero Avenue, San Francisco CA 94110. E-mail: msiobal@sfghsom.ucsf.edu.




Fig. 1. Effects of systemic vasodilation (from intravenous administration) versus selective pulmonary vasodilation (from inhaled administration) of prostacyclins (prostaglandin E₁ [PGE₁] and prostaglandin I₂ [PGI₂]). Systemic vasodilation affects all vascular beds, thereby decreasing mean arterial blood pressure and worsening oxygenation by increasing blood flow to poorly ventilated alveoli. Inhaled vasodilators selectively dilate pulmonary capillaries in alveoli that are well ventilated, thus reducing pulmonary artery pressure while improving ventilation/perfusion matching and oxygenation. However, "spillover" of inhaled drug to poorly ventilated alveoli and into the systemic circulation worsens shunt fraction and systemic hemodynamics. Q̇ = perfusion. V̇ = ventilation. MAP = mean arterial pressure. PAP = pulmonary artery pressure. PVR = pulmonary vascular resistance. SVR = systemic vascular resistance. Q̇_s/Q̇_t = shunt fraction.

hospital's drug formulary for the FDA-approved use in infants. Nonapproved and "off label" use of INO was prohibited because of its high cost and because it does not benefit outcomes in adults.⁷ At the request of the departments of anesthesia, pulmonary critical care, and respiratory care services, the pharmacy and therapeutics committee also added aerosolized prostacyclin (prostaglandin I₂ [PGI₂], brand name Flolan) to the hospital's drug formulary, based on evidence that inhaled prostacyclin benefits pulmonary hypertension, right-heart failure, and hypoxemic respiratory failure. We then developed a delivery system for aerosolized prostacyclin that allows easy dose calculation and adjustment,⁸ perioperatively, intraoperatively, and in the intensive care unit.

The present review describes the inhalable prostacyclins (PGI₂ and prostaglandin E₁ [PGE₁]) and their potential benefits for pulmonary hypertension, right-heart failure, and hypoxemia from acute respiratory distress syndrome (ARDS). I will also discuss the prostacyclins' potential benefits in decreasing platelet-aggregation and compare PGI₂ and INO with regard to duration of action, adverse effects, toxicology, methods of delivery during mechanical ventilation, problems associated with continuous aerosolization, and efficacy.

Prostacyclins Available for Aerosolization

Prostaglandin I₂

PGI₂ is a naturally occurring prostaglandin, a potent vasodilator, and an effective inhibitor of platelet aggregation, with an in vivo half-life of approximately 3–6 min. PGI₂ is FDA-approved for pulmonary hypertension via continuous intravenous infusion. PGI₂ must be reconstituted with a specific sterile diluent. Once reconstituted it is stable for 8 hours at room temperature, for 48 hours with refrigeration, and must be discarded after those time limits. The reconstituted solution of PGI₂ has a pH of 10.2–10.8 and is increasingly unstable at a lower pH. PGI₂ is photosensitive and must be protected from direct sunlight.⁹

Walmrath et al¹⁰ were first to publish data that directly compared the short-term use of aerosolized PGI₂ to INO in 16 mechanically ventilated ARDS patients. PGI₂ and INO were individually titrated in random order to maximize arterial oxygenation. Both aerosolized PGI₂ (at 7.5 ± 2.5 ng/kg/min, range 1.5–34 ng/kg/min) and INO (at 17.8 ± 2.7 ppm, range 2–40 ppm) improved oxygenation, reduced pulmonary shunt fraction (Q̇_s/Q̇_t), and lowered pulmonary artery pressure and PVR, with similar efficacy profiles (Fig. 2).

Iloprost

Iloprost, a stable analogue of PGI₂, is soluble in saline and has a plasma half-life of 20–30 min.¹¹ Inhaled iloprost, INO, and inhaled PGI₂ have similar pulmonary and hemodynamic effects in treating pulmonary hypertension.^{12–14} Hooper et al¹⁴ evaluated the short-term use of




Fig. 2. Comparison of aerosolized prostacyclin (prostaglandin I₂ [PGI₂]) (7.5 ± 2.5 ng/kg/min) and inhaled nitric oxide (NO) (17.8 ± 2.7 ppm) titrated to maximum oxygenation improvement in 16 patients with acute respiratory distress syndrome. PVR = pulmonary vascular resistance. P_aO₂/F_iO₂ = ratio of P_aO₂ to fraction of inspired oxygen ratio. MPAP = mean pulmonary artery pressure. Q̇_s/Q̇_t = shunt fraction. * p < 0.05. † p < 0.001. (Adapted from Reference 10.)




Fig. 3. Comparison of the effects of inhaled nitric oxide (INO at 40 ppm for 15 min) and a single dose of aerosolized iloprost (14–17 μ g over 15 min, approximately 15 ng/kg/min) in 35 patients with primary pulmonary hypertension. Values are mean \pm SD: A: Mean systemic arterial pressure (open squares) and pulmonary artery pressure (closed circles). B: Systemic vascular resistance (open squares) and pulmonary vascular resistance (open circles). C: Cardiac output. (Adapted from Reference 14.)

INO (at 40 ppm) versus a single dose of aerosolized iloprost (approximately 15 ng/kg/min for 15 min) in 35 patients with severe pulmonary hypertension (mean pulmonary artery pressure $59\text{--}60 \pm 11$ mm Hg). Inhaled iloprost had a greater effect in decreasing mean pulmonary artery pressure and PVR and increasing cardiac output and P_{aO_2} . However, inhaled iloprost significantly reduced both mean systemic arterial pressure and systemic vascular resistance (SVR) (Fig. 3), whereas INO did not. That effect on systemic vasomotor tone is presumably due to spillover into the systemic circulation, which is directly related to the longer duration of action (up to 2 h),^{12–14} which makes intermittent aerosolized iloprost a potential treatment for chronic pulmonary hypertension in ambulatory patients.¹⁵

In Europe inhaled iloprost has been studied in a large randomized, placebo-controlled trial,¹⁶ 2 small randomized comparisons to INO,^{17,18} several other observational studies,^{19–21} and case reports.^{22–25} Unfortunately, iloprost is not available in the United States.

Prostaglandin E₁

PGE₁ (generic name alprostadil, brand name Prostin VR Pediatric) is another naturally occurring prostaglandin; it causes vasodilation, inhibits platelet aggregation, and stimulates intestinal and uterine smooth muscle. Intravenous PGE₁ is used in neonates with congenital heart defects to maintain the patency of the ductus arteriosus until corrective surgery can be performed.²⁶ When administered intravenously PGE₁ is rapidly distributed and metabolized, has an estimated half-life of 5–10 min, and 70–80% of it is removed via the pulmonary vascular bed with a single pass through the lungs. Ampules (1 mL) of PGE₁ must be stored refrigerated. Contact of undiluted solution with the plastic sidewalls of volumetric infusion chambers must be avoided. PGE₁ diluted with saline or dextrose must be discarded after 24 hours.²⁷ Published reports of experience with inhaled PGE₁ is limited. In animals inhaled PGE₁ appeared to be less effective in reducing pulmonary hypertension during pharmacologically induced pulmonary vasoconstriction than was inhaled PGI₂ or INO.^{28,29} In 10 adult ARDS patients INO and both intravenous and aerosolized PGE₁ decreased mean pulmonary artery pressure and PVR and increased right-ventricular ejection fraction.³⁰ With PGE₁ inhalation requires a smaller mean dose than intravenous infusion (10 ± 1 vs 12 ± 2 ng/kg/min) for a similar effect in lowering pulmonary artery pressure. However, in contrast to inhaled PGE₁ and INO, intravenous infusion of PGE₁ causes systemic vasodilation, which lowers mean systemic arterial pressure and SVR and worsens arterial oxygenation and \dot{Q}_s/\dot{Q}_t (Fig. 4).

Indications and Evidence for Aerosolized Prostacyclin

The indications for use of aerosolized prostacyclins parallel the indications for INO, which include treatment of primary and secondary pulmonary hypertension, cardiac-surgery-associated pulmonary hypertension and right-ventricle failure, lung-transplantation-related reperfusion injury, liver transplantation that results in portopulmonary hypertension, hypoxemia due to single-lung ventilation or ARDS, and sickle cell disease.^{31,32} In numerous case reports and observational trials aerosolized prostacyclins have been effective for all of the above indications,^{1,12–25,30,33–39} though not for sickle-cell related vaso-occlusive crisis (acute chest syndrome).

AEROSOLIZED PROSTACYCLINS




Fig. 4. Hemodynamic and oxygen gas exchange variables during inhalation of aerosolized PGE₁ (10 ± 1 ng/kg/min) and inhaled nitric oxide (INO at 7 ± 1 ppm) in 10 adult patients with acute respiratory distress syndrome. A: Mean pulmonary artery pressure (PAP). B: Pulmonary vascular resistance (PVR). C: Right-ventricular ejection fraction (RVEF). D: Cardiac index. E: Systemic arterial pressure (SAP). F: Systemic vascular resistance (SVR). G: P_aO₂. H: Pulmonary shunt fraction (Q_s/Q_t). Panels E through H show the systemic hemodynamics and oxygenation effects of the nonselective vasodilation caused by intravenous infusion of PGE₁. (Data adapted from Reference 30.)

Pulmonary Hypertension

Inhaled vasodilators benefit acute and chronic pulmonary hypertension.^{1,33,40} Hache et al³³ retrospectively reviewed 37 patients who received inhaled PGI₂ over a 1-year

period. Twenty-two patients received aerosol boluses, 4 received continuous aerosolization, 9 received a combination of aerosol boluses and continuous aerosolization, 1 received aerosol treatment via face mask prior to intubation, and 1 received direct intratracheal boluses prior to

nebulization. Of the 27 patients who had pulmonary artery pressure monitoring, there was a significant decrease in mean pulmonary artery pressure, from 34.9 ± 11.8 to 32.1 ± 11.8 mm Hg ($p = 0.0017$), and the best response was 27.5 ± 11.1 mm Hg ($p < 0,0001$).

Oslichewski et al¹² tested the short-term effects of 15 min of INO (at 10–28 ppm), PGI₂ (53–115 ng/kg/min), and iloprost (9–22 ng/kg/min) based on an estimated average ideal body weight of 65 kg in 6 patients suffering severe pulmonary hypertension. PGI₂ and its long-acting analog, iloprost, had identical efficacy profiles. PGI₂ lowered pulmonary artery pressure 18%, whereas INO lowered pulmonary artery pressure only 10%; PGI₂ lowered PVR 41%, whereas PGI₂ lowered PVR 28%. Ten minutes of aerosolized PGI₂ (10 μ g/mL) was also compared to INO (40 ppm) in evaluating heart-transplant patients who were suffering elevated pulmonary artery pressure and PVR.³⁴ Inhaled PGI₂ and INO both reduced mean pulmonary artery pressure 7%. PGI₂ lowered PVR 49%, whereas INO lowered pulmonary artery pressure 43%. PGI₂ increased cardiac output 11%, whereas INO caused no change in cardiac output.

Aerosolized prostacyclins have been used with pre-term and term infants and with older pediatric patients who had pulmonary hypertension from prematurity, meconium aspiration, and congenital heart disease.^{25,41,42} In those case reports inhaled PGI₂ and iloprost used intraoperatively or in the intensive care unit reduced pulmonary artery pressure and improved oxygenation. Kelly et al⁴³ used inhaled PGI₂ to treat 4 hypoxemic term infants who had persistent pulmonary hypertension refractory to INO and found improvements in both P_{aO₂} (57 ± 6 to 100 ± 27 mm Hg, $p = 0.06$) and oxygenation index (29 ± 5 to 19 ± 7 , $p < 0.05$).

The development of pulmonary hypertension in patients undergoing coronary artery bypass grafting with cardiopulmonary bypass predicts higher mortality, perioperative myocardial infarction, and stroke.⁴⁴ The benefits of intraoperative and perioperative inhaled vasodilators to treat pulmonary hypertension is well documented in the anesthesia literature.^{33,35,40} Five patients treated intraoperatively with aerosolized PGI₂ (calculated average dose 35 ng/kg/min) during surgery had a small but significant decrease in pulmonary artery pressure (7%, $p < 0.03$) and a 35% reduction in PVR ($p < 0.004$).³⁵ Lowson et al³⁶ used a dose of 50 ng/kg/min of inhaled PGI₂ and reported pulmonary artery pressure reductions of 44% and 31% during 2 attempts to wean a patient from cardiopulmonary bypass following mitral and aortic valve replacement. Because of the technical complexity, cost, and potential toxicity of INO, inhaled PGI₂ is an effective alternative inhalable vasodilator for perioperative, intraoperative, and intensive care use.

Right Ventricular Failure

The pulmonary circulation is normally a low-pressure, low-resistance circuit that is highly distensible with recruitable vessels that can accommodate large increases in cardiac output. Acutely or chronically elevated pulmonary artery pressure increases PVR and right-ventricle afterload (the resistance the right ventricle pumps against) and results in a progressive inability of the right ventricle to sustain its flow output (decreased right ventricular stroke volume and right-ventricular ejection fraction). This eventually leads to elevated right-ventricle end-diastolic volume, right-ventricle hypertrophy, right-ventricle failure,⁴⁰ and in extreme cases can lead to left-ventricular pump failure in critically ill patients^{45,46} and higher mortality.^{47–50} Current state-of-the-art pulmonary-artery catheters allow continuous monitoring of right-ventricular ejection fraction, right-ventricle end-diastolic volume, and cardiac output,^{51–53} and thus allow close monitoring of the effects of inhaled pulmonary vasodilators on RV function.

Reducing right-ventricle afterload by decreasing PVR and pulmonary artery pressure is an important goal in the management of acute right-heart failure.^{32,35,54,55} Case reports by Schroeder et al³⁵ demonstrate the effectiveness of inhaled PGI₂ for altering right-ventricle function in cardiopulmonary patients. Patients treated with aerosolized PGI₂ (calculated average dose 35 ng/kg/min) for right-ventricle failure during cardiac and abdominal surgery had a 26% increase in cardiac index ($p < 0.003$) and a 35% decrease in PVR ($p < 0.004$). With 9 patients suffering pulmonary hypertension and right-ventricle failure, Haraldsson et al⁵⁶ studied the effects of inhaled PGI₂ after cardiac surgery and heart transplantation. Aerosolized PGI₂ (10 μ g/mL via continuous nebulization) improved right-ventricle performance and reduced PVR by 29%.

Inhaled PGI₂ is as effective as INO in lowering PVR, pulmonary artery pressure, and right-ventricle afterload. Because PGI₂ has a longer half-life and provides longer vasodilation than INO, it is a more potent inhaled vasodilator for pulmonary hypertension and right-ventricle failure. In dose-comparison trials the pulmonary artery pressure^{10,57,58} and PVR^{10,34,58} reductions were greater with PGI₂ than with INO (see Figs. 2 and 5).

Acute Respiratory Distress Syndrome

ARDS is characterized by severe hypoxemia from intrapulmonary shunting, areas of low \dot{V}/\dot{Q} ,⁵⁹ and pulmonary hypertension from elevated PVR.⁶⁰ Pulmonary hypertension develops early in ARDS,⁶¹ and the primary causes of the pulmonary hypertension are mechanical obstruction of the pulmonary microcirculation by microthromboemboli (composed of platelets and leukocytes) and hypoxic pulmonary vasoconstriction from alveolar and




Fig. 5. Dose-response curves comparing the effects of aerosolized prostacyclin (1, 10, and 25 ng/kg/min) and inhaled nitric oxide (INO at 1, 4, and 8 ppm) in 8 patients with acute respiratory distress syndrome. MPAP = mean pulmonary artery pressure. PVR = pulmonary vascular resistance. (Adapted from Reference 58.)

interstitial edema triggered by inflammation mediators.⁶¹⁻⁶³ Prostacyclin's antithrombotic and platelet-disaggregation effects may help prevent obstruction of pulmonary micro-circulation (endarteritis obliterans, inflammation, and fibrous tissue formation of the arterial inner wall) commonly seen postmortem in ARDS patients.² Pulmonary hypertension and the consequent right-ventricle dysfunction in ARDS patients predict higher mortality⁶¹⁻⁶⁴ and correlate with the severity of lung injury.^{61,63-65} Villar et al⁶⁴ found that among 225 patients suffering acute respiratory failure, 70 who had hemodynamic or pulmonary instability monitored via pulmonary-artery catheter had higher mortality (79%, 30/38) when pulmonary hypertension was present (mean pulmonary artery pressure 29 ± 6 mm Hg) than those who did not have pulmonary hypertension (44% mortality [14/32], mean pulmonary artery pressure 15 ± 3 mm Hg) ($p < 0.01$). Thirty of the 38 patients who had pulmonary hypertension also met ARDS diagnosis criteria, and their mortality was 70% (21/30). The 21 ARDS patients who died had significantly higher PVR and lower cardiac index than patients who did not die ($p < 0.001$).

Walmrath et al⁶⁶ were the first to report the effects of aerosolized PGI₂ for ARDS. In 3 patients with severe ARDS inhaled PGI₂ (at doses between 17 and 50 ng/kg/min) improved the ratio of P_{aO₂} to fraction of inspired oxygen (F_{IO₂}) by 44% (120 ± 19 to 173 ± 18 mm Hg), reduced shunt fraction by improving \dot{V}/\dot{Q} matching, decreased pulmonary artery pressure from 40.3 ± 13.5 to 32.0 ± 3.8 mm Hg, and lowered PVR by 30%.

Van Heerden et al⁶⁷ found marked oxygenation improvements and reduced \dot{Q}_s/\dot{Q}_t in 2 hypoxemic ARDS patients who received aerosolized PGI₂ doses of 50 ng/kg/min. The efficacy of aerosolized PGI₂ versus INO in hypoxemic ARDS patients has been studied in short-term observational trials in pediatric⁶⁸ and adult patients.^{10,66,67} Both PGI₂ and INO improve oxygenation and reduce \dot{Q}_s/\dot{Q}_t (see Figs. 2 and 5). Doses as low as 10 ng/kg/min improve oxygenation (Fig. 6).^{8,10,68,69} Inhaled prostacyclins cause minimal systemic vasodilation, are anti-inflammatory, and inhibit platelet aggregation,^{70,71} so they may be effective against the severe refractory hypoxemia and pulmonary-hypertension-induced right-ventricle dysfunction in severe ARDS.^{72,73}




Fig. 6. Dose-response curves showing the effects of aerosolized prostacyclin (at 0, 10, 20, 30, 40, and 50 ng/kg/min) on ratio of P_{aO_2} to fraction of inspired oxygen (P_{aO_2}/F_{iO_2}), mean arterial pressure (MAP), alveolar-arterial oxygen difference ($P_{(A-a)O_2}$), mean pulmonary artery pressure (MPAP), shunt fraction (shunt %), and cardiac index in 9 patients with hypoxemia due to acute respiratory distress syndrome (ARDS). (From Reference 69, with permission.)

Other Potential Benefits

The longer half-life of PGI_2 and spillover into the systemic circulation were suggested as the cause of the improved splanchnic perfusion observed by Eichelbronner et al.⁷⁴ Sixteen patients with pulmonary hypertension and septic shock that required norepinephrine and/or epinephrine to maintain mean arterial pressure > 65 mm Hg were randomized to received INO (19 ± 10 ppm) or inhaled PGI_2 (18 ± 9 ng/kg/min) until mean pulmonary artery pressure decreased by 15% (35 ± 4 to 30 ± 4 , $p < 0.05$, and 34 ± 4 to 30 ± 3 , $p < 0.05$ respectively). Neither INO nor PGI_2 affected systemic hemodynamics, cardiac index, or right-ventricular function, but PGI_2 nonsignificantly increased the indocyanine-green dye clearance rate (an index of hepatic blood flow) (6.7 to 4.8 min), whereas INO did not. In addition, unlike INO, inhaled PGI_2 significantly increased gastric pH, from 7.26 ± 0.07 to 7.30 ± 0.05 ($p < 0.05$) and reduced the arterial-gastric mucosal P_{CO_2} gradient from 19 ± 6 to 15 ± 4 ($p < 0.05$), indicating

improved splanchnic perfusion. Eichelbronner et al also hypothesized that the PGI_2 platelet-aggregation inhibition, antithrombotic, and anti-inflammatory effects might have contributed to the observed improved splanchnic perfusion.

Prostacyclin also stimulates endothelial release of nitric oxide,⁷⁵ which suggests that prostacyclin may have an additive benefit when used in combination with other therapies.² This was confirmed in a study by Della Rocca et al,⁷⁶ which demonstrated that combined INO and inhaled PGI_2 reduced pulmonary artery pressure and improved P_{aO_2}/F_{iO_2} significantly more than INO alone.

Inhaled prostacyclins have also been administered in combination with both aerosolized and oral phosphodiesterase inhibitors in animal models of pulmonary hypertension^{77,78} and in human subjects with severe pulmonary hypertension.¹⁷ Ghofrani et al¹⁷ compared (1) INO (at 20 to 40 ppm), (2) the combination of oral sildenafil (12.5 and 50 mg) and inhaled iloprost, and (3) oral sildenafil alone, in random order, with 30 patients suffering severe pulmo-

nary hypertension (class IV by the classification system of the New York Heart Association). Combining oral sildenafil (both doses) and inhaled iloprost enhanced and prolonged the vasodilatory effect without affecting systemic arterial pressure or oxygenation. Patients who received 50 mg of sildenafil and inhaled iloprost had the greatest reduction in pulmonary vascular resistance, -44.2% compared with -14.1% for INO. The synergistic effect of combining other vasodilators with inhaled prostacyclins may be an effective strategy in treating severe pulmonary hypertension.

Adverse Effects and Toxicity

The potent vasodilation and platelet-aggregation-inhibition effects of PGI₂ make systemic hypotension and bleeding the most important potential adverse effects. The effective dose range established by dose-response trials^{10,58,66–69,79–81} is 5–50 ng/kg/min. No systemic hemodynamic effects have been reported with that dose range. Systemic hypotension was reported in one patient when the aerosolized dose exceeded 200 ng/kg/min³⁷ and in a series of 5 healthy male volunteers who inhaled approximately 700 ng/kg/min.⁴ Moreover, hypotension from an overdose of PGI₂ can be rapidly reversed with supportive therapy and discontinuation of PGI₂.⁹ Problems with bleeding have not been reported, but it would be reasonable to avoid aerosolized PGI₂ during active pulmonary hemorrhage.

The effect of intrapulmonary and systemic spillover of inhaled PGI₂ may help reduce pulmonary hypertension and improve splanchnic perfusion but simultaneously may adversely affect \dot{V}/\dot{Q} matching, oxygenation, and systemic hemodynamics (see Fig. 1). Among pneumonia patients with and without pulmonary fibrosis, higher doses of aerosolized PGI₂ (33.6 ± 12.0 vs 6.6 ± 3.0 ng/kg/min) were required to reduce mean pulmonary artery pressure in patients who had fibrosis.⁸¹ That higher dose significantly reduced P_{aO_2}/F_{IO_2} (from 73.8 ± 6.6 to 65.5 ± 6.8 mm Hg, $p < 0.05$), increased $\dot{Q}s/\dot{Q}t$ (from 44.7 ± 3.0 to $49.4 \pm 5.0\%$), decreased mean arterial pressure (from 80.3 ± 3.6 to 71.3 ± 4.7 mm Hg, not a significant difference), and reduced SVR (755 ± 120 to 701 ± 115 dyn·cm⁻⁵, not a significant difference).

In another study, mean arterial pressure and SVR were insignificantly lowered (3.7% and 10.0%, respectively) by aerosolized PGI₂ (compared to INO) in heart transplant patients who had pulmonary hypertension.³⁴ And in yet another study, intraoperative aerosolized PGI₂ decreased mean arterial pressure by 5.2% and significantly reduced SVR (23.3%).³⁵ The adverse effects of spillover on $\dot{Q}s/\dot{Q}t$ and systemic vasodilation are probably offset by the improvements in \dot{V}/\dot{Q} matching and right-ventricle function, but clinicians should be aware of this potential effect.

Common adverse effects during initial intravenous administration of PGI₂ are generally related to vasodilation and include (in order of frequency) flushing, headache, nausea, vomiting, hypotension, anxiety, chest pain, and dizziness.⁹ The degree and frequency of those adverse effects has not been established and none have been reported from inhaled PGI₂.

As with INO the accidental or intentional abrupt withdrawal of inhaled PGI₂ could cause rebound pulmonary vasoconstriction, acute \dot{V}/\dot{Q} mismatch, hypoxemia, pulmonary hypertension, and right-ventricle failure, though the potential for those adverse effects may be less with aerosolized PGI₂ than with INO, because PGI₂ has a longer half-life and duration of action (20–25 min vs 5 min for INO).¹⁰ The potential for rebound may also be reduced by weaning slowly.

Unlike INO, prostacyclin has no known toxic effects or toxic metabolites. However, reconstituted PGI₂ solution has a very alkaline pH (10.2–10.8)⁹ that may act as an irritant when inhaled. Habler et al^{82,83} found no evidence of substantial acute pulmonary toxicity, lung tissue damage, or increase in bleeding time in lambs that received doses and volumes of PGI₂ solution equivalent to what would be administered to humans (28 ng/kg/min). Conversely, development of mild acute sterile tracheitis (polymorphonuclear leukocyte infiltration) was observed in piglets that received aerosolized PGI₂ at doses and volumes approximately 9 times the maximum that would be given to an adult human patient.⁸⁴ The importance of those findings is unclear, but we have had experience with one patient who received aerosolized PGI₂ via face mask and developed a severe coughing episode during a cardiac catheterization procedure. We have administered aerosolized PGI₂ therapy for as long as 4.8 days to intubated patients⁸ and have observed no adverse effects, but we advise caution when administering this extremely alkaline aerosol to patients with reactive airways disease.

Aerosol Delivery Methods

Aerosol administration of drugs is a very inefficient method of delivery in that as little as 3% of the nominal dose deposits in the lungs.⁸⁵ Aerosolized PGI₂ therapy requires continuous aerosolization usually during mechanical ventilation over an extended period, from several hours to as long as several days. It is essential to select an appropriate nebulizer and delivery method to promote alveolar deposition and ensure a stable dose administration.

Alveolar deposition is favored when the aerosol particles are in the range of 1–2 μm in diameter.⁸⁶ Therefore nebulizers that produce a small median particle size are optimal for aerosolized PGI₂ therapy. Various jet nebulizers, with an average particle size of 3.4 μm (range 2.1–5.2 μm) and an average driving flow of 6 L/min




Fig. 7. Syringe pump delivery system for aerosolized prostacyclin. A: Syringe pump with 50-mL syringe of diluted prostacyclin. B: Connector to endotracheal tube. C: Jet nebulizer. D: Ventilator. E: Flow meter for air-oxygen mixture (6 L/min). (From Reference 69, with permission.)

(range 4–12 L/min), can effectively administer aerosolized PGI₂.^{10,12,35,38,57,58,66,67,69,81} Delivery via ultrasonic nebulizers with particle sizes of 2.5 μm and 4.0 μm has also been reported.^{68,80}

The evaporation of solvent during both jet and ultrasonic nebulization gradually concentrates the drug solution in the nebulizer.⁸⁷ Steckel and Eskandar found a greater drug-concentration effect with an ultrasonic nebulizer than with a jet nebulizer (48% vs 13% increase respectively).⁸⁸ Ultrasonic nebulization generates heat and the medication may reach 40–55°C,^{89–91} which results in higher evaporation and concentration effects. Reconstituted PGI₂ in the specified diluent solution remains stable at room temperature (15–25°C) for 8 hours.⁹ It has not been determined whether the heating from ultrasonic nebulization substantially affects the potency of PGI₂. Recent modifications in ultrasonic nebulizer design have reduced aerosol particle size and the evaporative concentration effect,^{92,93} and only ultrasonic nebulizers with those operating characteristics should be used for prolonged aerosolized PGI₂ therapy.

Jet nebulizers have several inherent properties that affect dose delivery during prolonged continuous nebulization. Aerosol output and particle size from a jet nebulizer partly depend on the flow rate of the gas powering the nebulizer⁹⁴ and the pneumatic pressure during operation.^{95,96} Jet nebulizers operating at different flows can have a large range of aerosol volume output and particle size, so there is the potential of large variability in dose delivery.⁹⁷ Operating a jet nebulizer from an external flow source alters tidal volume, accuracy of exhaled tidal volume measurement, ventilator circuit pressures, patient-initiated triggering, and F_{IO₂} if an external oxygen blender is not used. Use of a built-in nebulizer-driver function on a ventilator also affects the delivered dose over time,⁹⁸ with various ventilation settings, such as respiratory rate and inspiratory time. Variability in the performance of individual nebulizers of the same model and manufacturer can also influence




Fig. 8. A delivery system for aerosolized prostacyclin. A drug solution concentration of 30 $\mu\text{g}/\text{mL}$ enables administration of dose ranges of 10–50 ng/kg/min to patients with predicted body weights between 40 and 100 kg.

dose delivery.^{99,100} Furthermore, during operation the volume loss from a jet nebulizer is the sum of the aerosol produced plus the evaporative loss of solvent. Because water vapor does not carry any drug, the amount of drug emitted from a jet nebulizer does not equal the amount calculated based on the volume loss from the nebulizer.^{8,96} Therefore, during continuous jet nebulization the initial aerosolized dose is always lower than the intended dose. Solvent evaporation concentrates drug in the nebulizer solution, which alters the delivered dose over time.⁸

Delivery methods for aerosolized PGI₂ therapy have been described vaguely^{58,80} and have had restrictive dose titration capability. The delivery system used by Van Heerden et al⁵⁷ incorporates a syringe pump and jet nebulizer with 6 L/min flow (Fig. 7). Setting or adjusting dose delivery with that system requires mixing the appropriate drug concentration in the syringe and adjusting the pump infusion rate to obtain the desired dose, based on the patient's body weight. The delivery method we developed at my institution uses a dual infusion pump system with a low-flow (approximately 2 L/min) jet nebulizer that en-

Table 1. Evidence Ranking of Studies Cited in This Report That Support the Use of Aerosolized Prostacyclin for Treating Pulmonary Hypertension and/or Hypoxemia

Evidence Level*	PGI ₂ References	PGE ₁ References	Iloprost References
I: Large randomized controlled trials	none	none	16
II: Small randomized controlled trials	none	none	none
III: Non-randomized, concurrent cohort controls	none	none	none
IV: Non-randomized, historical controls	10, 13, 34, 57, 58, 74, 80† 12, 38, 56, 66-69, 76, 79, 81	30†	13, 14, 17, 18† 12, 15, 19, 20
V: Case series reports	8, 33, 35-37, 39, 41-43	none	22-25

* Evidence Level I is highest (strongest evidence). Level V is lowest.

PGI₂ = prostaglandin I₂.
PGE₁ = prostaglandin E₁.

† Comparisons to inhaled nitric oxide.
(Adapted from References 102 and 103.)

Table 2. Grades of Recommendation for Aerosolized Prostacyclin

Grade of Recommendation*	Prostacyclin Type
Grade A: Supported by at least 2 Level I investigations	none
Grade B: Supported by at least 1 Level I investigation	iloprost
Grade C: Supported by at least 2 Level II investigations	none
Grade E: Supported by at least 1 Level III investigation	none
Grade F: Supported by Level IV or Level V investigations	PGI ₂ , PGE ₁

* Grade A is highest (strongest evidence). Grade F is lowest.

PGI₂ = prostaglandin I₂.
PGE₁ = prostaglandin E₁.
(Adapted from Reference 105.)

ables the administration of a wide range of doses and dose titrations over a wide range of patient body weights, using a single drug solution concentration (Fig. 8).⁸

Certain new (and currently available) electronic nebulizers do not generate heat but do produce a constant volume output of small aerosol particles, without an external flow source.¹⁰¹ That eliminates interference with ventilator function and the problems associated with solvent evaporation and dose variability. Unfortunately, these new nebulizers are designed for intermittent use and are not FDA approved for continuous nebulization.

Efficacy of Aerosolized Prostacyclin

Examination of the hierarchy of evidence in the studies cited in this report, using standard methodology,¹⁰²⁻¹⁰⁴ reveals primarily low levels of supporting evidence for aerosolized prostacyclins (Table 1). The efficacy of aerosolized iloprost is supported by one large, randomized, placebo-controlled trial of the treatment of severe pulmonary hypertension.¹⁶ In the established hierarchy of evidence-grading¹⁰⁵ the evidence supporting aerosolized

iloprost qualifies for a grade B recommendation (Table 2). In contrast, the results from predominately short-term, uncontrolled studies of aerosolized PGI₂ and PGE₁ merit only a grade F (the lowest) recommendation. A recent Cochrane Review of prostacyclin for pulmonary hypertension¹⁰⁶ recognized the potential benefit of aerosolized iloprost, compared to other treatments, but the efficacy of aerosolized PGI₂ and PGE₁ was not assessed.

The evidence is clear that inhaled prostacyclins have effects similar to INO in improving oxygenation in ARDS and reducing pulmonary hypertension.^{10,13,14,17,18,30,34,57,58,74,80} Also, similar to INO, only around 60% of patients respond to aerosolized PGI₂.^{8,10,79} Numerous early trials of INO for hypoxemia and pulmonary hypertension in ARDS also looked promising, but 4 level-1 randomized, controlled clinical studies failed to show positive differences in important patient outcomes.¹⁰⁷ However, the similarities of prostacyclins and INO should not disqualify prostacyclin as a treatment option. Randomized, controlled trials are needed to determine the efficacy of inhaled prostacyclin.

Summary

Prostacyclins are potent vasodilators, they inhibit platelet aggregation, and they are anti-inflammatory. Aerosolized prostacyclins reduce pulmonary artery pressure, improve right-heart function, and increase arterial oxygenation by improving \dot{V}/\dot{Q} matching. Their longer duration of action and spillover into the systemic circulation may prove to have additional benefits over other inhaled vasodilators. There are no known serious toxic effects or toxic metabolites associated with aerosolized prostacyclins. Current nebulizers that are designed for continuous aerosol generation suffer variable dose delivery. The efficacy of aerosolized prostacyclin therapy has not been proven in large, randomized, controlled studies.

REFERENCES

- Lowson SM. Inhaled alternatives to nitric oxide. *Anesthesiology* 2002;96(6):1504–1513.
- Wetzel RC. Aerosolized prostacyclin: in search of the ideal pulmonary vasodilator (editorial). *Anesthesiology* 1995;82(6):1315–1317.
- Szczeklik A, Gryglewski RJ, Nizankowski E, Nizankowski R, Musial J. Pulmonary and anti-platelet effects of intravenous and inhaled prostacyclin in man. *Prostaglandins* 1978;16(5):651–660.
- Burghuber OC, Silberbauer K, Haber P, Sinzinger H, Elliott M, Leithner C. Pulmonary and antiaggregatory effects of prostacyclin after inhalation and intravenous infusion. *Respiration* 1984;45(4):450–454.
- Hardy C, Robinson C, Lewis RA, Tattersfield AE, Holgate ST. Airway and cardiovascular responses to inhaled prostacyclin in normal and asthmatic subjects. *Am Rev Respir Dis* 1985;131(1):18–21.
- Ignarro LJ, Buga GM, Wood KS, Byrns RE, Chaudhuri G. Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proc Natl Acad Sci USA* 1987;84(24):9265–9269.
- Dellinger RP, Zimmerman JL, Taylor RW, Straube RC, Hauser DL, Criner GJ, et al. Effects of inhaled nitric oxide in patients with acute respiratory distress syndrome: results of a randomized phase II trial. Inhaled Nitric Oxide in ARDS Study Group. *Crit Care Med* 1998;26(1):15–23.
- Siobal MS, Kallett RH, Pittet JF, Warnecke EL, Kraemer RW, Venkayya RV, Tang JF. Description and evaluation of a delivery system for aerosolized prostacyclin. *Respir Care* 2003;48(8):742–753.
- Glaxo Wellcome SmithKline, Flolan Product Information. July 2001.
- Walmrath D, Schneider T, Schermuly R, Olschewski H, Grimminger F, Seeger W. Direct comparison of inhaled nitric oxide and aerosolized prostacyclin in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1996;153(3):991–996.
- Schering, Ilomedin Product DataSheet. September 2001.
- Olschewski H, Walmrath D, Schermuly R, Ghofrani HA, Grimminger F, Seeger W. Aerosolized prostacyclin and iloprost in severe pulmonary hypertension. *Ann Intern Med* 1996;124(9):820–824.
- Olschewski H, Ghofrani HA, Walmrath D, Schermuly R, Temmesfeld-Wollbrück B, Grimminger F, Seeger W. Inhaled prostacyclin and iloprost in severe pulmonary hypertension secondary to lung fibrosis. *Am J Respir Crit Care Med* 1999;160(2):600–607.
- Hoepfer MM, Olschewski H, Ghofrani HA, Wilkins H, Winkler J, Borst MM, et al. A comparison of the acute hemodynamic effects of inhaled nitric oxide and aerosolized iloprost in primary pulmonary hypertension. *J Am Coll Cardiol* 2000;35(1):176–182.
- Hoepfer MM, Schwarze M, Ehlerdin S, Adler-Schuermeyer A, Spiekerkoetter E, Niedermeyer J, et al. Long-term treatment of primary pulmonary hypertension with aerosolized iloprost, a prostacyclin analogue. *N Engl J Med* 2000;342(25):1866–1870.
- Olschewski H, Simonneau G, Galie N, Higenbottam T, Naeije R, Rubin LJ, et al. Inhaled iloprost for severe pulmonary hypertension. *N Engl J Med* 2002;347(5):322–329.
- Ghofrani HA, Wiedemann R, Rose F, Olschewski H, Schermuly RT, Weissmann N, et al. Combination therapy with oral sildenafil and inhaled iloprost for severe pulmonary hypertension. *Ann Intern Med* 2002;136(7):515–522.
- Sablotzki A, Hentschel T, Gruenig E, Schubert S, Friedrich I, Muhling J, et al. Hemodynamic effects of inhaled aerosolized iloprost and inhaled nitric oxide in heart transplant candidates with elevated pulmonary vascular resistance. *Eur J Cardiothorac Surg* 2002;22(5):746–752.
- Kramm T, Eberle B, Krummenauer F, Guth S, Oelert H, Mayer E. Inhaled iloprost in patients with chronic thromboembolic pulmonary hypertension: effects before and after pulmonary thromboendarterectomy. *Ann Thorac Surg* 2003;76(3):711–718.
- Theodoraki K, Rellia P, Thanopoulos A, Tsourelis L, Zarkalis D, Sfyraakis P, Antoniou T. Inhaled iloprost controls pulmonary hypertension after cardiopulmonary bypass. *Can J Anaesth* 2002;49(9):963–967.
- Machherndl S, Kneussl M, Baumgartner H, Schneider B, Petkov V, Schenk P, Lang IM. Long-term treatment of pulmonary hypertension with aerosolized iloprost. *Eur Respir J* 2001;17(1):8–13.
- Muller M, Scholz S, Kwapisz M, Akinturk H, Thul J, Hempelmann G. Use of inhaled iloprost in a case of pulmonary hypertension during pediatric congenital heart surgery. *Anesthesiology* 2003;99(3):743–744.
- Rex S, Busch T, Vettelschoss M, de Rossi L, Rossaint R, Buhre W. Intraoperative management of severe pulmonary hypertension during cardiac surgery with inhaled iloprost. *Anesthesiology* 2003;99(3):745–747.
- Wittwer T, Pethig K, Struber M, Hoepfer M, Harringer W, Haverich A, Franke U, Wahlers T. Aerosolized iloprost for severe pulmonary hypertension as a bridge to heart transplantation. *Ann Thorac Surg* 2001;71(3):1004–1006.
- Langer F, Wendler O, Wilhelm W, Tscholl D, Schafers HJ. Treatment of a case of acute right heart failure by inhalation of iloprost, a long-acting prostacyclin analogue. *Eur J Anaesthesiol* 2001;18(11):770–773.
- Pharmacia and Upjohn, Prostin VR Pediatric, Package Insert, August 2000.
- RxMed.com. Pharmaceutical information: prostin VR. Accessed April 7, 2004.
- Walmrath D, Schermuly R, Pilch J, Grimminger F, Seeger W. Effects of inhaled versus intravenous vasodilators in experimental pulmonary hypertension. *Eur Respir J* 1997;10(5):1084–1092.
- Kleen M, Habler O, Hofstetter C, Pusch R, Mueller M, Welte M, Zwissler B. Efficacy of inhaled prostanoids in experimental pulmonary hypertension. *Crit Care Med* 1998;26(6):1103–1109. Erratum in: *Crit Care Med* 1998;26(7):1294.
- Putensen C, Hormann C, Kleinsasser A, Putensen-Himmer G. Cardiopulmonary effects of aerosolized prostaglandin E₁ and nitric oxide inhalation in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1998;157(6 Pt 1):1743–1747.
- Stuedel W, Hurford WE, Zapol WM. Inhaled nitric oxide: basic biology and clinical applications. *Anesthesiology* 1999;91(4):1090–1121.
- Kleen M, Zwissler B. Intraoperative use of inhaled vasodilators: are there indications? *Curr Opin Anaesth* 2002;15(1):79–83.
- Hache M, Denault AY, Belisle S, Couture P, Babin D, Tetrault F, Guimond JG. Inhaled prostacyclin (PGI₂) is an effective addition to the treatment of pulmonary hypertension and hypoxemia in the operating room and intensive care unit. *Can J Anaesth* 2001;48(9):924–929.
- Haraldsson A, Kieler-Jensen N, Nathorst-Westfelt U, Bergh CH, Ricksten SE. Comparison of inhaled nitric oxide and inhaled aerosolized prostacyclin in the evaluation of heart transplant candidate with elevated pulmonary vascular resistance. *Chest* 1998;114(3):780–786.
- Schroeder RA, Wood GL, Plotkin JS, Kuo PC. Intraoperative use of inhaled PGI₂ for acute pulmonary hypertension and right ventricular failure. *Anesth Analg* 2000;91(2):291–295.
- Lowson SM, Doctor A, Walsh BK, Doorley PA. Inhaled prostacyclin for the treatment of pulmonary hypertension after cardiac surgery. *Crit Care Med* 2002;30(12):2762–2764.
- Webb SAR, Stott S, van Heerden PV. The use of inhaled aerosolized prostacyclin (IAP) in the treatment of pulmonary hypertension

- secondary to pulmonary embolism. *Intensive Care Med* 1996;22(4):353–355.
38. Della Rocca G, Coccia C, Costa MG, Pompei L, Di Marco P, Vizza CD, et al. Inhaled aerosolized prostacyclin and pulmonary hypertension during anesthesia for lung transplantation. *Transplant Proc* 2001;33(1–2):1634–1636.
 39. Schroeder RA, Rafii AA, Plotkin JS, Johnson LB, Rustgi VK, Kuo PC. Use of aerosolized inhaled epoprostenol in the treatment of portopulmonary hypertension. *Transplantation* 2000;70(3):548–550.
 40. Fischer LG, Van Aken H, Burkle H. Management of pulmonary hypertension: physiological and pharmacological considerations for anesthesiologists. *Anesth Analg* 2003;96(6):1603–1616.
 41. Zwissler B, Rank N, Jaenicke U, Schurle B, Welte M, Reichart B, et al. Selective pulmonary vasodilation by inhaled prostacyclin in a newborn with congenital heart disease and cardiopulmonary bypass. *Anesthesiology* 1995;82(6):1512–1516.
 42. Soditt V, Aring C, Groneck P. Improvement of oxygenation induced by aerosolized prostacyclin in a preterm infant with persistent pulmonary hypertension of the newborn. *Intensive Care Med* 1997;23(12):1275–1278.
 43. Kelly LK, Porta NFM, Goodman DM, Carrol CL, Steinhorn RH. Inhaled prostacyclin for term infants with persistent pulmonary hypertension refractory to inhaled nitric oxide. *J Pediatr* 2002;141(6):830–832.
 44. Reich DL, Bodian CA, Krol M, Kuroda M, Osinski T, Thys DM. Intraoperative hemodynamic predictors of mortality, stroke, and myocardial infarction after coronary artery bypass surgery. *Anesth Analg* 1999;89(4):814–822.
 45. Sibbald WJ, Driedger AA. Right ventricular function in acute disease states: pathophysiologic considerations. *Crit Care Med* 1983;11(5):339–345.
 46. Sibbald WJ, Driedger AA, Myers ML, Short AI, Wells GA. Biventricular function in the adult respiratory distress syndrome. *Chest* 1983;84(2):126–134.
 47. Eddy AC, Rice CL, Anardi DM. Right ventricular dysfunction in multiple trauma victims. *Am J Surg* 1988;155(5):712–715.
 48. Hurford WE, Zapol WM. The right ventricle and critical illness: a review of anatomy, physiology, and clinical evaluation of its function. *Intensive Care Med* 1988;14 Suppl 2:448–457.
 49. Dhainaut JF, Lanore JJ, de Gournay JM, Huyghebaert MF, Brunet F, Villemant D, Monsallier JF. Right ventricular dysfunction in patients with septic shock. *Intensive Care Med* 1988;14 Suppl 2:488–491.
 50. Eddy AC, Rice CL. The right ventricle: an emerging concern in the multiply injured patient. *J Crit Care* 1989;4(1):58–66.
 51. Vincent JL, Thirion M, Brimiouille S, Lejeune P, Kahn RJ. Thermodilution measurement of right ventricular ejection fraction with a modified pulmonary artery catheter. *Intensive Care Med* 1986;12(1):33–38.
 52. Dhainaut JF, Brunet F, Monsallier JF, Villemant D, Devaux JY, Konno M, et al. Bedside evaluation of right ventricular performance using a rapid computerized thermodilution method. *Crit Care Med* 1987;15(2):148–152.
 53. Nelson LD. The new pulmonary arterial catheters: right ventricular ejection fraction and continuous cardiac output. *Crit Care Clin* 1996;12(4):795–818.
 54. Stobierska-Dzierzek B, Awad H, Michler RE. The evolving management of acute right-sided heart failure in cardiac transplant recipients. *J Am Coll Cardiol* 2001;38(4):923–931.
 55. Zwissler B. [Acute right heart failure: etiology, pathophysiology, diagnosis, therapy.] *Anaesthesist* 2000;49(9):788–808. Article in German.
 56. Haraldsson A, Kieler-Jensen N, Ricksten SE. Inhaled prostacyclin for treatment of pulmonary hypertension after cardiac surgery or heart transplantation: a pharmacodynamic study. *J Cardiothorac Vasc Anesth* 1996;10(7):864–868.
 57. Van Heerden PV, Blythe D, Webb SAR. Inhaled aerosolized prostacyclin and nitric oxide as selective pulmonary vasodilators in ARDS—a pilot study. *Anaesth Intensive Care* 1996;24(5):564–568.
 58. Zwissler B, Kemming G, Habler O, Kleen M, Merkel M, Haller M, et al. Inhaled prostacyclin (PGI₂) versus inhaled nitric oxide in adult respiratory distress syndrome. *Am J Respir Crit Care Med* 1996;154(6 Pt 1):1671–1677.
 59. Dantzer DR, Brook CJ, Dehart P, Lynch JP, Weg JG. Ventilation-perfusion distributions in the adult respiratory distress syndrome. *Am Rev Respir Dis* 1979;120(5):1039–1052.
 60. Zapol Wm, Snider MT. Pulmonary hypertension in severe acute respiratory failure. *N Engl J Med* 1977;29(9):476–480.
 61. Moloney ED, Evans TW. Pathophysiology and pharmacological treatment of pulmonary hypertension in acute respiratory distress syndrome. *Eur Respir J* 2003;21(4):720–727.
 62. Spapen H, Vincken W. Pulmonary arterial hypertension in sepsis and the adult respiratory distress syndrome. *Acta Clin Belg* 1992;47(1):30–41.
 63. Romand JA, Donald FA, Suter PM. Cardiopulmonary interactions in acute lung injury: clinical and prognostic importance of pulmonary hypertension. *New Horiz* 1994;2(4):457–462.
 64. Villar J, Blazquez MA, Lubillo S, Quintana J, Manzano JL. Pulmonary hypertension in acute respiratory failure. *Crit Care Med* 1989;17(6):523–526.
 65. Leeman M. Pulmonary hypertension in acute respiratory distress syndrome. *Monaldi Arch Chest Dis* 1999;54(2):146–149.
 66. Walmrath D, Schneider T, Pilch J, Grimminger F, Seeger W. Aerosolized prostacyclin in adult respiratory distress syndrome. *Lancet* 1993;342(8877):961–962.
 67. Van Heerden PV, Webb SAR, Hee G, Corkeron M, Thompson WR. Inhaled aerosolized prostacyclin as a selective pulmonary vasodilator for the treatment of severe hypoxemia. *Anaesth Intens Care* 1996;24(1):87–90.
 68. Pappert D, Busch T, Gerlach H, Lewandowski K, Radermacher P, Rossaint R. Aerosolized prostacyclin versus inhaled nitric oxide in children with severe acute respiratory distress syndrome. *Anesthesiology* 1995;82(6):1507–1511.
 69. van Heerden PV, Barden A, Michalopoulos N, Bulsara MK, Roberts BL. Dose-response to inhaled aerosolized prostacyclin for hypoxemia due to ARDS. *Chest* 2000;117(3):819–827.
 70. Vane JR, Anggard EE, Botting RM. Regulatory functions of the vascular endothelium. *N Engl J Med* 1990;323(1):27–36.
 71. Tateson JE, Moncada S, Vane JR. Effects of prostacyclin (PGX) on cyclic AMP concentrations in human platelets. *Prostaglandins* 1977;13(3):389–397.
 72. Grimminger R, Rose F, Ghofrani HA, Schermuly RT, Weissmann N, Olschewski H, et al. [Inhalative strategies for improvement of pulmonary hemodynamics and gas exchange in sepsis and severe pulmonary hypertension.] *Z Kardiol* 2000;89(6):477–484. Article in German.
 73. Kaisers U, Busch T, Deja M, Donaubauber B, Falke KJ. Selective pulmonary vasodilation in acute respiratory distress syndrome. *Crit Care Med* 2003;31(4 Suppl):S337–342.
 74. Eichelbronner O, Reinelt H, Wiedeck H, Mezody M, Rossaint R, Georgieff M, Radermacher P. Aerosolized prostacyclin and inhaled nitric oxide in septic shock—different effects on splanchnic oxygenation? *Intensive Care Med* 1996;22(9):880–887.
 75. Shimokawa H, Flavahan NA, Lorenz RR, Vanhoutte PM. Prostacyclin releases endothelium-derived relaxing factor and potentiates its action in coronary arteries of the pig. *Br J Pharmacol* 1988;95(4):1197–1203.

76. Della Rocca G, Coccia C, Pompei L, Ruberto F, Venuta F, De Giacomo T, Pietropaoli P. Hemodynamic and oxygenation changes of combined therapy with inhaled nitric oxide and inhaled aerosolized prostacyclin. *J Cardiothorac Vasc Anesth* 2001; 15(2):224–227.
77. Schermuly RT, Krupnik E, Tenor H, Schudt C, Weissmann N, Rose F, et al. Coaerosolization of phosphodiesterase inhibitors markedly enhances the pulmonary vasodilatory response to inhaled iloprost in experimental pulmonary hypertension: maintenance of lung selectivity. *Am J Respir Crit Care Med* 2001; 164(9):1694–1700.
78. Schermuly RT, Roehl A, Weissmann N, Ghofrani HA, Leuchte H, Grimminger F, et al. Combination of nonspecific PDE inhibitors with inhaled prostacyclin in experimental pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol* 2001;281(6):L1361–L1368.
79. Bein T, Metz C, Keyl C, Sendtner E, Pfeifer M. Cardiovascular and pulmonary effects of aerosolized prostacyclin administration in severe respiratory failure using a ventilator nebulization system. *J Cardiovasc Pharmacol* 1996;27(4):583–586.
80. Domenighetti G, Stricker H, Waldspuehl B. Nebulized prostacyclin (PGI₂) in acute respiratory distress syndrome: impact of primary (pulmonary injury) and secondary (extrapulmonary injury) disease on gas exchange response. *Crit Care Med* 2001; 29(1):57–62.
81. Walmrath D, Schneider T, Pilch J, Schermuly R, Grimminger F, Seeger W. Effects of aerosolized prostacyclin in severe pneumonia: impact of fibrosis. *Am J Respir Crit Care Med* 1995;151(3 Pt 1):724–730.
82. Habler O, Kleen M, Zwissler B, Pusch R, Welte M, Vogelmeier C, et al. Inhalation of prostacyclin (PGI₂) for eight hours does not produce signs of acute pulmonary toxicity in healthy lambs. *Intensive Care Med* 1996;22(5):426–433.
83. Habler O, Kleen M, Takenaka S, Leiderer R, Pusch R, Welte M, Zwissler B, Messmer K. Eight hours' inhalation of prostacyclin (PGI₂) in healthy lambs: effects on tracheal, bronchial, and alveolar morphology. *Intensive Care Med* 1996;22(11):1232–1238.
84. van Heerden PV, Caterina P, Filion P, Spagnolo DV, Gibbs NM. Pulmonary toxicity of inhaled aerosolized prostacyclin therapy: an observational study. *Anaesth Intensive Care* 2000;28(2):161–166.
85. MacIntyre NR, Silver RM, Miller CW, Schuler F, Coleman RE. Aerosol delivery in intubated, mechanically ventilated patients. *Crit Care Med* 1985;13(2):81–84.
86. Stuart BO. Deposition of inhaled aerosols. *Arch Intern Med* 1973; 131(1):60–73.
87. O'Callaghan C, Barry PW. The science of nebulised drug delivery. *Thorax* 1997;52 Suppl 2:S31–S44.
88. Steckel H, Eskandar F. Factors affecting aerosol performance during nebulization with jet and ultrasonic nebulizers. *Eur J Pharm Sci* 2003;19(5):443–455.
89. Phipps PR, Gonda I. Droplets produced by medical nebulizers: some factors affecting their size and solute concentration. *Chest* 1990;97(6):1327–1332.
90. Niven RW, Ip AY, Mittelmann S, Prestrelski SJ, Arakawa T. Some factors associated with the ultrasonic nebulization of proteins. *Pharm Res* 1995;12(1):53–59.
91. Servo i Ventilator System V. 1.4 Users Manual; U. S. Addition: 229.
92. Katial RK, Reisner C, Buchmeier A, Bartelson BB, Nelson HS. Comparison of three commercial ultrasonic nebulizers. *Ann Allergy Asthma Immunol* 2000;84(2):255–261.
93. Flament MP, Leterme P, Gayot A. Study of the technological parameters of ultrasonic nebulization. *Drug Dev Ind Pharm* 2001; 27(7):643–649.
94. Hess D, Fisher D, Williams P, Pooler S, Kacmarek RM. Medication nebulizer performance: effects of diluent volume, nebulizer flow, and nebulizer brand. *Chest* 1996;110(2):498–505.
95. Clay MM, Pavia D, Newman SP, Clarke SW. Factors influencing the size distribution of aerosols from jet nebulisers. *Thorax* 1983;38(10):755–759.
96. Raabe OG, Wong TM, Wong GB, Roxburgh JW, Piper SD, Lee JIC. Continuous nebulization therapy for asthma with aerosols of β_2 agonists. *Ann Allergy Asthma Immunol* 1998;80(6):499–508.
97. Rau JL. Design principles of liquid nebulization devices currently in use. *Respir Care* 2002;47(11):1257–1275; discussion 1275–1278.
98. Hughes JM, Saez J. Effects of nebulizer mode and position in a mechanical ventilator circuit on dose efficiency. *Respir Care* 1987; 32(12):1131–1135.
99. Hollie MC, Malone RA, Skufca RM, Nelson HS. Extreme variability in aerosol output of the DeVilbiss 646 jet nebulizer. *Chest* 1991;100(5):1339–1344.
100. Alvine GF, Rodgers P, Fitzsimmons KM, Ahrens RC. Disposable jet nebulizers. How reliable are they? *Chest* 1992;101(2):316–319.
101. Dhand R. Nebulizers that use a vibrating mesh or plate with multiple apertures to generate aerosol. *Respir Care* 2002;47(12):1406–1416; discussion 1416–1418.
102. Sacket DL. Rules of evidence and clinical recommendations on the use of antithrombotic agents. *Chest* 1986;89(2 Suppl):2S–3S.
103. Cochrane and Systematic Reviews. Levels of evidence for health-care interventions. Available at: <http://www.cochrane.org/consumers/sysrev.htm#levelsofevidence>. Accessed February 24, 2004.
104. Montori VM, Guyatt GH. What is evidenced-based medicine and why it should be practiced? *Respir Care* 2001;46(11):1201–1214.
105. Pulmonary Artery Catheter Consensus conference: consensus statement. *Crit Care Med* 1997;25(6):910–925.
106. Paramothayan NS, Lasserson TJ, Wells AU, Walters EH. Prostacyclin for pulmonary hypertension. *Cochrane Database Syst Rev* 2003;(2):CD002994.
107. Kopp R, Kuhlen R, Max M, Rossaint R. Evidence-based medicine in the therapy of the acute respiratory distress syndrome. *Intensive Care Med* 2002;28(3):244–255.