

**Summary**

Grey Heron (*Ardea cinerea*) swallows a juvenile Great Crested Grebe (*Podiceps cristatus*)

In the special protected area „Kühkopf-Knoblochsaue“ (Hesse) we observed a Grey Heron (*Ardea cinerea*) swallowing a juvenile Great Crested Grebe (*Podiceps cristatus*), probably being dead already before. Based on the literature it is one of the largest birds caught by a Grey Heron so far.

**Literatur**

Bauer, K., & U. Glutz von Blotzheim (1966): Handbuch der Vögel Mitteleuropas, Bd. 1: Gaviiformes – Phoenicopteriformes. Akademische Verlagsgesellschaft, Frankfurt a. M. \* Creutz, G. (1981): Der Graureiher. Neue Brehm Bücherei, Bd. 530, A. Ziems Verlag, Wittenberg-Lutherstadt. \* Hormann, M. & M. Korn (1994): Bestandsentwicklung ausgewählter, gefährdeter Vogelarten in Hessen 1990 bis 1993. – Ergebnisse der Indikatorartenauswertung. – Vogel und Umwelt 8 (3): 147–159. \* Murphy, C. M. (1976): Grey Herons eating Water Rails. Brit. Birds 69: 369. \* Voisin, C. (1991): The Herons of Europe. Poyser, London.

Josef Kreuziger und Ernst-Ludwig Achenbach

Anschrift der Verfasser: JK, Gartenstraße 17, D-64673 Zwingenberg, ELA, Brandenburgerstr. 6,  
D-64297 Darmstadt.

---

## Responses of the Crane (*Grus grus*) to potential predators in traditional wintering areas

References to the anti-predatory behaviour of Cranes are very scarce. CRAMP & SIMMONS (1980) have referred to a sudden grouping of birds on ground, with subdued alarm calls when standing birds are attacked by White-tailed Eagle (*Haliaeetus albicilla*) in breeding areas. Unlike MUÑOZ-PULIDO et al. (1993) pointed out that the normal response of Cranes when they discover Golden Eagles (*Aquila chrysaetos*) or Griffon Vultures (*Gyps fulvus*) is to fly high in closely-packed groups.

In the present paper we show the responses of Cranes to seven potential predators in the Serena region, SW of Spain. This is a traditional wintering area of the Western route of this species (SÁNCHEZ et al. 1993). During the winter periods of 1994–1995 and 1995–1996 a total of 20 encounters of Crane flocks with potential predators were recorded (Table 1). Taking into account that

Table 1: Antipredator behaviour in the Crane in the presence of different predator species. No response means that Cranes did not change their activities when the predator appeared.

Tab. 1: Feindabwehrverhalten von Kranichen gegenüber verschiedenen Beutegreifernarten, „No response“ bedeutet, daß die Vögel keine Aktivitätsänderungen bei Herannahen des jeweiligen Beutegreifers zeigten.

Species	Flight	Grouping	No response
Golden Eagle ( <i>Aquila chrysaetos</i> )	6	1	1
Bonelli's Eagle ( <i>Hieraetus fasciatus</i> )	1	–	–
Griffon Vulture ( <i>Gyps fulvus</i> )	–	1	1
Buzzard ( <i>Buteo buteo</i> )	1	–	1
Red Kite ( <i>Milvus milvus</i> )	–	–	4
Raven ( <i>Corvus corax</i> )	–	1	1
Red fox ( <i>Vulpes vulpes</i> )	–	1	–

we detected a change in the behaviour of Cranes in the presence of the Raven (*Corvus corax*) and in the presence of the Buzzard (*Buteo buteo*), these species were included as potential predators. The inclusion of the Fox (*Vulpes vulpes*) was based on existing reports of this species hunting cranes in North America (DRIESLIN & BENNET 1979).

The proportion of flight responses was higher when eagles (*Aquila chrysaetos* and *Hieraetus fasciatus*) hovered over the flocks of Cranes than when other predators appeared (Log-likelihood ratio Fisher test:  $G = 302.6$ ,  $p < 0.01$ ). Predation of Cranes by Golden Eagles has been described (THYOLLAY 1979, MUÑOZ-PULIDO et al. 1993, IVANOSKY 1995). Therefore the responses in the presence of this species and of Bonelli's Eagle could be the consequence of the Crane's recognition of potential predation by these species, and the response in this case would involve a greater expenditure of energy to decrease the chances of predation by increasing the confusion effect caused by the predator's difficulty in selecting its prey from a group in movement (PULLIAN & CARACO 1984).

The Griffon Vulture, contrary to the reported by MUÑOZ-PULIDO et al. (1993), the Red Fox and the Raven induced a grouping in Crane flocks. At the present only attacks on young chicks during the breeding period by *Vulpes* genus has been recorded (DRIESLIN & BENNET 1979), while egg predation by *Corvus* genus seem to occur in the presence of adults without eliciting response (CRAMP & SIMMONS 1980). Probably the grouping creates the saturation of predator and prevents preying on individuals (TURCHIN & KAREIVA 1989) from species that are potential predators during the breeding period, when doesn't exit Crane aggregation (CRAMP & SIMMONS 1980). Unlike the Red Kite (*Milvus milvus*) do not seem to have any predatory capacity on the Crane.

### Acknowledgements

We are specially grateful to D. Parejo by discussion on previous drafts. An anonymous referee made useful comments on an earlier draft of the manuscript.

### Zusammenfassung

Die Reaktionen in SW-Spanien (Region Serena) überwinternder Kraniche gegenüber sieben potentiellen Beutegreifern werden dargestellt. Drei verschiedene Verhaltensäußerungen beim Erscheinen von Prädatoren wurden festgestellt: Auffliegen, Bildung dichter Gruppen am Boden sowie unverändert fortgeführte Aktivitäten (keine Reaktion). Das Auffliegen wurde am ehesten von Stein- und Habichtsadlern, Gruppenbildung am Boden hingegen in einigen Fällen von Gänsegeier, Kolkrabe und Fuchs ausgelöst. Lediglich Rotmilane blieben völlig unbeachtet.

### References

- Cramp, S., & K. E. L. Simmons (1980): The birds of the Western Palearctic Vol. 2. Oxford University press. \*Drieslein, R. L., & A. J. Bennett (1979): Red Fox predation on Greater Sandhill Crane chicks. Wilson Bulletin 91: 132–133. \*Ivanovsky, V. V. (1995): The Common Crane in the system of trophic relationships of predatory birds in Bielorussia. In: Crane research and protection in Europe (H. Prange editor). Martin-Luther-Universität, Halle-Wittenberg. \*Muñoz-Pulido, R. Alonso, J. C. & J. A. Alonso (1993). Common Crane (*Grus grus*) killed by Golden Eagle (*Aquila chrysaetos*). Vogelwarte 37: 78–79. \*Pulliam, H. R. & T. Caraco (1984). Living in groups: is there an optimal group size? In Behavioural Ecology and Evolutionary Approach. Edited by J. R. Krebs & N. B. Davies. Blackwell Scientific Publications. \*Sánchez, J. M., A. Sánchez, A. Fernández & A. Muñoz (1993): La Grulla común (*Grus grus*) en Extremadura status y relación con el uso del suelo. Edita Grupo de Investigación en Conservación, Área de Biología Animal, Universidad de Extremadura. \*Thiollay, J. M. (1979): La migration des Grues à travers l'Himalaya et la prédation par les Aigles Royaux. Alauda 47: 83–92. \*Turchin, P., & P. Kareiva (1989). Aggregation in *Aphis varians*: an effective strategy for reducing predation risk. Ecology 70: 1008–1016.

Jesús M. Avilés, Juan. M. Sánchez and Francisco J. Medina

Addresses of the authors. Grupo de Investigación en Conservación, Área de Biología Animal, Universidad de Extremadura, 06071. Badajoz, Spain.

# ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Vogelwarte - Zeitschrift für Vogelkunde](#)

Jahr/Year: 1997/98

Band/Volume: [39\\_1998](#)

Autor(en)/Author(s): Aviles Jesus M., Sanchez Juan M.

Artikel/Article: [Responses of the Crane \(\*Grus grus\*\) to potential predators in traditional wintering areas 302-303](#)